Powerful pollinators

Encouraging insect pollinators in rural and urban landscapes



Pollinators are an essential component of healthy, biodiverse landscapes and provide critical pollination services to native flora and agriculture production across the country.

This guide provides information on ways to encourage a diverse range of insect pollinators across all properties, and includes a planting calendar to help select plants to support diverse pollinators throughout the year.



The power of pollinators

Pollinators – mostly insects, but also birds and mammals – assist the production of seeds and fruit in many plant species by visiting flowers in search of food (nectar and/or pollen). Whilst foraging they transfer pollen from one flower to another, facilitating fertilisation, which results in fruits and seeds.

Honey bees, native bees and other native insects like hoverflies, wasps and butterflies provide essential pollination services for native plants, pastures, crops, fruits and vegetables.

Pollinators and food security

Without insect pollinators, the quantity and diversity of food grown for humans in contemporary agricultural systems would be severely restricted. Many of the food crops we eat, as well as pasture and fodder crops, benefit from pollination by insects.

Pollinator-dependent crops include lucerne, faba beans, apples and vegetables, as well as many crops grown for seed production, such as canola.

The quantity and diversity of insect pollinators are key drivers of production as they influence both crop yields and quality. Under-pollination results in smaller and misshapen fruit or seed that isn't viable.

Grazing enterprises can also suffer from a reduction in the abundance or diversity of pollinators, due to the role these insects play in the persistence of nitrogen-fixing pasture legumes such as clover.

A diverse and healthy community of pollinators generally provides more effective and consistent pollination than relying on any single species.



Native vegetation supports pollinators by providing food and nesting sites. Nearby crops and pastures will benefit from the increased abundance and diversity of pollinators in the landscape.

Insect populations are in decline worldwide due to land clearing, intensive or monocultural agriculture, pesticide use, pollution, colony disease, increased urbanisation and climate change. Low pollinator numbers mean not all flowers are pollinated, leading to low fruit or seed set. This in turn reduces fruit and vegetable harvest yields, and decreases food supply.



Under-pollination results in smaller, misshapen fruit such as this strawberry.

Healthy ecosystems

Pollinators are both essential to, and depend upon, healthy ecosystems. A growing human population and increasing demand for food puts pressure on ecosystems, while declining ecosystem function will in turn negatively impact food production.

Insect pollinators are a prime example of this — without healthy ecosystems and the presence of patches of native vegetation to support insect populations, pollination will decline. This will threaten both crop productivity and the persistence of native, pollinator-dependent flowering plants.

Pollinators require habitat that contains year-round food sources, breeding resources and nesting sites. The presence of pollinator habitat adjacent to food crops has been shown to improve food production by enabling a greater variety and number of pollinators to persist year-round, providing pollination services when required.

Turn to the centre of this brochure for a guide to planting for pollinators.

Diapause or diet? Where are the insects?

Many insect pollinators undergo a diapause during colder winter months. Diapause is a period of suspended development during unfavourable environmental conditions, and during this period insect pollinators do not need flowers. Birds and other small mammals will continue to benefit from available pollen and nectar during this time.

If there are low numbers of insect pollinators in your local area, it is important to determine whether this is because of diapause, or because of an inadequate availability of nectar and pollen, creating a 'food desert' where insect pollinators cannot survive.

There are still many unknowns about insect pollinators in Australia. Take part in Australian Pollinator Week or in the annual Australian Pollinator Count to learn more about pollinators in your area – visit: **AustralianPollinatorWeek.org.au** and **AustralianPollinatorCount.au**

Encouraging pollinators on your property

Create pollination reservoirs

Pollination reservoirs are areas of native plant species that provide floral resources for pollinators. They can be new plantings or existing habitat, such as shelterbelts or remnant vegetation. A high diversity of plant species is essential to provide nectar, pollen and nesting sites throughout the year. Pollination reservoirs need to be close enough to crops to ensure that pollinators can fly easily to them.

Use existing habitat

Protect and improve existing habitat where possible. Roadsides, shelterbelts, dam margins, woodlands, grasslands, rocky areas, river and creek edges can all be important pollinator-attracting areas, bringing valuable pollination services to your property.

Native vegetation stands provide habitat for pollinators, and special attention should be paid to enhance and protect these areas.

Get to know your local flora

Each property and region will have distinct populations of insects, based on the plants and climate. Identifying and understanding the insects in your area will help you develop better plantings.

The plants growing in nearby bush will be well suited to the climate and soils in your region. Local community groups and specialist native nurseries can provide useful information and usually produce local plant species.

Plant trees, shrubs and groundcovers

Planting a variety of species of groundcovers, shrubs and trees on your property will further attract pollinators to your area. Use a combination of direct seed sowing and planting tube stock to establish new vegetation. Initial watering and protection from grazing will improve the success rate of young plants. Wildflowers, including our native pea species, are excellent at attracting a diverse range of native pollinators.

Connectivity counts

Insect pollinators benefit from greater connectivity of habitat in a landscape, which allows them to forage over a wider radius and increase in numbers in a local area. Encourage neighbours and other landholders to plant for pollinators and create connections across your landscape.

Utilise ecotones

Ecotones are the margins between two different habitats. Ecotones often contain a more diverse mixture of pollinator species because they are inhabited by pollinators from both habitats. Protect and utilise ecotones such as the transition zones between woodland and grassland, or heath and shrubland, to create highly diverse floral and insect communities.

Amplify the flower signal

Plants have evolved large flowers or clusters of smaller flowers which attract more pollinator visits. Large, colourful and diverse plantings attract more pollinators. Ideally, plant in groups that contain different vegetation layers – combine a species-rich mixture of wildflowers, groundcovers, herbs, lilies, rushes, climbers, shrubs and trees.

Plant for the future

When establishing pollinator habitat, consider including species that are indigenous to your area and can tolerate increasingly warmer and drier (or wetter) conditions, to improve resilience to climate change. Rehabilitate weedy areas into managed pollination reservoirs by introducing lots of flowering plant diversity. Be careful not to plant invasive or listed weeds, and look for suitable replacements.

Double the crop value

Plants that are pollinator-attracting may be commercially viable crop species in their own right and can be used to diversify farm production. Bush foods such as Wirilda (*Acacia*) seed, Pig Face, Sweet Apple Berry and many more are in high demand for use in fresh and manufactured products. Native plant seed is also needed for revegetation projects. Farmers can also support beekeepers by hosting beehives to increase pollinator numbers on a farm.

Reduce chemical use

Insecticides, fungicides and herbicides all affect the health of bees, bee colonies and native pollinators. Herbicides can impact pollinators by reducing the availability and diversity of flora and removing vegetation that helps support insect life. Some herbicides can also harm the beneficial microbes in the insect gut. In many circumstances, beneficial insects will, in healthy numbers, help control pest insects, ultimately reducing the need for insecticide use.

When chemical pest control is unavoidable, select products that are least harmful for pollinators and apply insecticides in the evening or at night when pollinators are not active.

Always use according to directions, especially for withholding periods, and notify beekeepers a few days before spraying chemicals so beehives can be safely relocated away from harm.

Be a citizen scientist and do some detective work to discover local pollinators on your property. Visit **inaturalist.ala.org.au** to be involved.

Safeguard the bees? The best way to 'save the bees' and protect our pollinators is to create an abundance of diverse habitat – from the ground up! There is much interest in keeping a beehive to promote pollinators, but there are serious legal and biosecurity responsibilities that must be considered, and that the introduction of a beehive does not displace existing native pollinators and insects. Be a friend of pollinators and say it with flowers! A guide to planting for pollinators for the Grampians region, Victoria

Healthy populations of insect pollinators are important for crop yields, orchard production and thriving native vegetation.

This planting guide will help you choose plant species to attract and keep pollinators on your property throughout the year.

The Greater Grampians bioregion in western Victoria is centred on the Grampians Mountain Range (Gariwerd). • flower colour and flowering season The Grampians is a dramatic landscape dominated by parallel ranges of resistant sandstone with valleys that have been cut either in soft shales or deeply weathered granites. The steep escarpments and gentle back slopes offer diversity of habitats for the region's The coloured bars indicate the flowering rich flora, comprising a third Victorian indigenous plants. Reaching over 1100 m, the Grampians has much higher rainfall than the surrounding plains. Following the winter rains, the Grampians is famous differ between regions and seasons, for its spectacular spring wildflower displays. In part, these occur due to the incredible diversity of habitats in the region, including wet forest, montane heath, grassy woodland and sandplain heath. The species cropped in the Grampians region vary from north to south due to a steep rainfall gradient.

The plants listed in this Guide will help supply rewards to pollinators, with an emphasis on species that are indigenous can't source these plants at your local and suited to local climates.

The eucalypt species in the chart have been selected as high quality honey production species. Most eucalypts do not flower every year, so choosing diverse species will help create continuously flowering habitat.



The pollinator plant list

To create pollinator-attracting plantings, use the Guide to choose a selection of plants with a variety of flower colours, different growth habits and a range of flowering seasons.

For each species, the planting Guide lists:

- life-form/'habit' (climber, herb, shrub or tree) and height (m).
- the vegetation type in which they naturally occur
- growth requirements (sun/shade, moist/dry)
- insect groups that may visit each plant and the floral reward (pollen and/or nectar).

months for each species. Darker shading denotes the peak flowering period, with a lighter shading for non-peak flowering months. Flowering dates may particularly for non-peak times, if your local climate is consistently warmer or cooler than average, with earlier or later flowering.

Sourcing plants

Most of the plant species listed are available from retail or wholesale nurseries or native plant growers, and local environment groups. If you garden centre, or indigenous nursery, ask them to contact the local nursery suppliers and plant growers listed online. See the reverse of the Guide for details.



Lifeform	Common name	Scientific name	Family	Vegetation type	Height	Flower colour		Flowering	Aspect	Soil moisture	Pollinator Pollen	Nector	Visitation by pollinator			y pollinator	e Mothe	Reatles	Flies
Crop plants							عر				rollen		dive bees noney	bees noven		sps Dufferfile		Deenes	Tiles
Herb	Canola	Brassica napus	Brassicaceae	Broadacre Cropping	1.5 m	Yellow	•	Final State Stat	Full sun	Well-drained	•	•	• •	•		•			•
Herb	Lucerne	Medicago sativa	Fabaceae	Broadacre Cropping	< 0.75 m	Purple			Full sun	Well-drained	•	•	• •						
Herb	Clover	Trifolium repens	Fabaceae	Pasture / Fodder	< 0.3 m	White		F F F F F F F F F F F F F F F F F F F	Full sun	Well-drained	•	•	• •						
Herb	Faba Beans Olive	Vicia faba	Fabaceae	Broadacre Cropping	<1m	White, Brown			-ull sun	Well-drained	•	•	• •						
Indiaenous plants	Olive	Olea europaea	Oleacede	Orchara	< 12 m	white			-uli sun	well-drained		•	• •	•					_
Prostrate	Running Postman	Kennedia prostrata	Fabaceae	Forest, Woodland, Heath	< 0.1 m	Red			Sun to shade	Well-drained	•	•	•						
Herb	Pale Vanilla Lily	Arthropodium milleflorum	Asparagaceae	Woodland, Grassy Woodland, Grassland	< 1.3 m	White, Pink			Sun to semi-shade	Well-drained	•*		•						
Herb	Chocolate Lily	Arthropodium strictum	Asparagaceae	Grassland, Grassy Woodland	<1m	Purple			Sun to semi-shade	Well-drained	•*		•						
Herb	Bulbine Lily	Bulbine bulbosa	Asphodelaceae	Grassland, Grassy Woodland	< 0.5 m	Yellow			Sun to semi-shade	Winter moist	•		• •						
Herb	Milkmaids Blue Stars	Burchardia umbellata	Colchicaceae	Grassland, Grassy Woodland	< 0.6 m	White			Sun to semi-shade	Winter moist	•	•	• •	•					•
Herb	Billy-Buttons	Craspedia variabilis	Asteraceae	Grassland, Grassy Woodland, Forest	< 0.25 m	Yellow			Sun, semi-shade, shade	Well-drained	•	•	• •	•			_		
Herb	Black-Anther Flax Lily	Dianella revoluta	Asphodelaceae	Heathland, Woodland, Forest	< 1.2 m	Purple			Sun to semi-shade	Well-drained	•*		• •						
Herb	Swamp Goodenia	Goodenia humilis	Goodeniaceae	Swamp & Lake Margins	< 0.2 m	Yellow			Full sun to shade	Moist	•	•	• •	•					
Herb	Tall Lobelia	Lobelia gibbosa	Campanulaceae	Heathland, Woodland, Forest	< 0.6 m	Purple			Sun to semi-shade	Well-drained	•	•	• •						
Herb	Yam Daisy	Microseris walteri	Asteraceae	Grassland, Grassy Woodland	< 0.5 m	Yellow			Sun to semi-shade	Winter moist	•	•	• •	•					
Herb	Long Purple-Flag	Patersonia occidentalis		Woodland, Heathland	< 0.55 m	Purple			-ull sun to shade	Poorly drained	•		• •						
Herb	Grass Triggerplant	Stylidium araminifolium	Stylidiaceae	Forest, Woodland, Heathland	< 0.0 m	Pink			Sun to semi-shade	Well-drained	•	•	• •	•		•	-		•
Herb	Nodding Blue Lily	Stypandra glauca	Asphodelaceae	Forest, Woodland, Heathland	< 1.5 m	Purple	•		Sun to semi-shade	Well-drained	•*	-	•						
Herb	Common Fringe Lily	Thysanotus tuberosus	Asparagaceae	Heathland, Woodland, Forest	< 0.4 m	Purple			Sun to semi-shade	Well-drained	•*		• •						
Herb	Hairy Annual-Bluebell	Wahlenbergia gracilenta	Campanulaceae	Woodland	< 0.5 m	Blue			Sun to semi-shade	Well-drained	•	•	• •						
Herb	Early Nancy	Wurmbea dioica	Colchicaceae	Woodland, Open Forest	< 0.2 m	Cream, Purple			Sun to semi-shade	Winter moist	•	•	• •	•				-	•
Herb	Austral Grasstree	Xanthorrhoea australis	Asphodelaceae	Heathland, Heathy Woodland, Forest	< 3 m	Yellow			Sun to semi-shade	Well-drained	•	•	• •	•	•		•	•	•
Sedae	Variable Sword-Sedae	Lepidosperma laterale	Cyperaceae	Heathy Woodland, Open Forest	<1m	Brown			Sun to semi-shade	Well-drained	•	•	• •						
Shrub	Golden Wattle	Acacia pycnantha	Fabaceae	Forest, Woodland	< 6 m	Yellow			Semi-shade to shade	Well-drained	•	•	• •	•					
Shrub	Juniper Wattle	Acacia ulicifolia	Fabaceae	Open Forest, Woodland, Heathland	< 2 m	Yellow	•		Sun to semi-shade	Well-drained	•		• •	•					
Shrub	Honey-Pots	Acrotriche serrulata	Ericaceae	Heathland, Heathy Woodland	< 0.3 m	White	0		Sun to semi-shade	Well-drained	•	•	•		•				•
Shrub	Brush Heath	Brachyloma ericoides	Ericaceae	Heath, Heathy Woodland	< 0.6 m	Pink			Sun to semi-shade	Well-drained	•	•	• •						
Shrub	Scarlet Bottlebrush	Callistemon rugulosus	Myrtaceae	Forest, Heath Along Creeks	< 3 m	Red			Sun to semi-shade	Moist	•	•	• •			•			
Shrub		Calytrix tetragona	Rutaceae	Forest Woodland Heathland	< 3 m	Red Green			Sun to semi-shade	Well-drained	•	•	• •	•			•	•	•
Shrub	Leafless Bitter-Pea	Daviesia brevifolia	Fabaceae	Forest, Woodland, Heathland	<1m	Red, Orange			Sun to semi-shade	Well-drained	•	•	• •						
Shrub	Gorse Bitter-Pea	Daviesia ulicifolia	Fabaceae	Forest, Woodland, Heathland	<1m	Yellow, Orange 😑	ŏ.		Sun to semi-shade	Well-drained	•	•	• •						
Shrub	Showy Parrot-Pea	Dillwynia sericea	Fabaceae	Heathland, Heathy Woodland	< 0.5 m	Red, Orange 🛛 🔴			Sun to semi-shade	Well-drained	•	•	• •						
Shrub	Common Heath	Epacris impressa	Ericaceae	Heath, Heathy Woodland, Forest	< 1.2 m	Pink, White	0		Full sun to shade	Well-drained	•	•	• •			•	•		
Shrub	Common Eutaxia	Eutaxia microphylla	Fabaceae	Woodland, Heathland	<1m	Yellow			Sun to semi-shade	Well-drained	•	•	• •						
Shrub	Hop Goodenia	Goodenia ovata	Goodeniaceae	Forest Woodland	< 0.5 m	Yellow			Full sun to shade	Well-drained	•	•							
Shrub	Holly Grevillea	Grevillea aquifolium	Proteaceae	Heathland, Woodland, Swampy Areas	< 2 m	Red	•		Sun to semi-shade	Well-drained	•	•	• •						
Shrub	, Bushy Needlewood	Hakea decurrens	Proteaceae	Open forest, Woodland, Heathland	< 3 m	Cream	$\overline{\mathbf{O}}$		Sun to semi-shade	Well-drained	•	•	• •	•					•
Shrub	Bundled Guinea-Flower	Hibbertia fasciculata	Dilleniaceae	Heathland, Heathy Woodland	< 0.5 m	Yellow	•	F	Full sun	Well-drained	•		• •	•				•	
Shrub	Silky Guinea-Flower	Hibbertia sericea	Dilleniaceae	Heathland, Heathy Woodland	<1m	Yellow			Sun to semi-shade	Well-drained	•		• •	•				•	
Shrub	Twiggy Guinea-Flower	Hibbertia virgata	Dilleniaceae	Heathland, Heathy Woodland	< 0.9 m	Yellow			Full sun	Well-drained	•		• •	•				•	
Shrub	Prickly ledfree Heath Teatree	Leptospermum continentale	Myrtaceae	Woodland, Heathland Heathland Heathy Woodland	< 2 m	White	$\frac{1}{2}$		Sun to semi-shade	Well-drained	•	•		•	_			•	
Shrub	Common Beard-Heath	Leucopogon virgatus	Ericaceae	Forest, Woodland, Heathland	< 0.6 m	White	$\overline{0}$		Full sun to shade	Well-drained	•	•	• •	•			•	•	•
Shrub	Scented Paperbark	Melaleuca squarrosa	Myrtaceae	Swamps	< 5 m	Yellow	Ŏ		Sun to semi-shade	Swampy	•	•	• •	•		•	•	•	•
Shrub	Prickly Broom-Heath	Monotoca scoparia	Ericaceae	Heath, Heathy Woodland	< 2 m	White	\bigcirc		Sun to semi-shade	Well-drained	•	•	• •	•		•	•	•	•
Shrub	Fringed Daisy-Bush	Olearia ciliata	Asteraceae	Heathland, Woodland, Forest	< 0.8 m	Purple			Sun to semi-shade	Well-drained	•	•	• •	•		•	_		
Shrub	Prickly Geebung	Persoonia juniperina	Proteaceae	Heathland, Heathy Woodland	<1m	Yellow			Sun to semi-shade	Well-drained	•	•	• •						
Shrub	Common Elat-Peg	Platylobium obtusanaulum	Fabaceae	Woodland Heathland	< 1.2 m	Yellow Red			Sun to semi-shade	Well-drained	•	•	• •			•	•		
Shrub	Victorian Christmas Bush	Prostanthera lasianthos	Lamiaceae	Forest, Moist Gullies	< 3 m	White, Pink			Semi-shade to shade	Moist	•	•	• •	•			•	•	•
Shrub	Rough Bush-Pea	Pultenaea scabra	Fabaceae	Moist Forest, Heathland	< 2 m	Yellow, Red			Sun to semi-shade	Well-drained	•	•	• •						
Shrub	Flame Heath	Stenanthera conostephioides	Ericaceae	Heath, Heathy Woodland	<1m	Red			Sun to semi-shade	Well-drained	•	•	• •						
Shrub	Pink-Bells	Tetratheca ciliata	Elaeocarpaceae	Forest, Woodland, Heathland	<1m	Pink		F F	Full sun to shade	Well-drained	•	•	• •						
Shrub	Grampians Thryptomene	Thryptomene calycina	Myrtaceae	Forest, Woodland, Heathland	< 2 m	White, Pink			Sun to semi-shade	Well-drained	•	•	• •	•			•	•	•
Tree	Buloke	Allocasuarina luehmannii	Casuarinaceae	woodland	< 15 m	Red			-uii sun Full sun	Well-drained		•	• •						
Tree	Black Wattle	Acacia meannsii	Fabaceae	Forest	< 10 m	Yellow			Semi-shade to shade	Well-drained	•	•		•					
Tree	Brown Stringybark	Eucalyptus baxteri	Myrtaceae	Forest, Woodland	< 40 m	Cream			Semi-shade	Moist	•	•	• •	•		•	•	•	•
Tree	River Red-Gum	Eucalyptus camaldulensis	Myrtaceae	Low-Lying Forest, Woodland, Floodplain	< 40 m	White	Õ		Semi-shade	Winter wet	•	•	•	•		•	•	•	•
Tree	Grampians Peppermint	Eucalyptus falciformis	Myrtaceae	Forest, Woodland	< 15m	White	0		Semi-shade	Well-drained	•	•	• •	•		•	•	•	•
Tree	Yellow-Gum	Eucalyptus leucoxylon	Myrtaceae	Woodland	< 25 m	White			Semi-shade	Well-drained	•	•	• •	•		•	•	•	•
Vine	Purple Coral Pea	Hardenbergia violacea	Fabaceae	Open Forest, Woodland	Vine	Purple			Sun to semi-shade	Well-drained	•	•	• •						
vine	Orange Bell-Climber	iviariantnus bignoniaceus	Pittosporaceae	vegetation Along Creeks	vine	Green, Orange 📒			berni-snade to shade	IVIOIST	•	•	•						

*Buzz Pollinated

Know your pollinators



European honey bees have two pairs of wings and long, segmented antennae. They are daytime-flying and feed on nectar and pollen. They are generalist pollinators and provide the bulk of pollination services for horticulture and crop plants. Honey bees and native bees are both essential to functioning ecosystems and food security in Australia.

Honey bees have become an important part of the Australian landscape. Honey bees live as colonies, and have a long history of coexistence with humans, including in domestic gardens.





Hoverfly (Family Syrobidae) (Family Syrobidae) **Australian native bees** comprise more than 2000 species, which provide essential pollination services. Native bees are generally solitary and live in nests in the ground or in hollow stems, old borer holes and other cracks and crevices, and some have evolved to pollinate particular native flowers through 'buzz pollination'. Although many Australian native bees are generalist foragers, some species have co-evolved with native plants and adapted to be the most effective pollinators of their flowers. Many native plant species, such as *Dianella* and *Grevillea* require specially adapted insects to access their nectar and enable the transfer of pollen to the stigma. Most native bees are solitary, but some species found in northern Australia (*Tetragonula* sp. and *Austroplebeia* sp.) are social bees and are used for commercial pollination of crops like macadamia nuts.

Fly species number up to 30,000 in Australia, and can be identified by having only one pair of flight wings. A second set of wings are modified into club-shaped paddles that allow flies to hover and stabilise their flight. Unlike bees and wasps, many flies (Brachycera) have very small, clubbed antennae at the front of their head. Flies, including blowflies, are often attracted to flowers that smell like carrion. Some flower-flies, have hairy bodies that easily collect pollen while they are feeding. Flies provide a range of services in the garden, including pollination, decomposition and predation.

Hoverflies are a type of fly, distinguishable by their large eyes, short antennae, bright black and yellow abdomen and their hovering flight behaviour. Adult hoverflies are nectar and pollen feeders. Hoverfly larvae feed on pests such as aphids, thrips and leafhoppers and are excellent biocontrol agents.



Beetles have hard outer wings that form their distinctive beetle shape. Their outer wings form a T-shape where they join at the top, unlike bugs where the outer wings make an X- or Y-shape. Some beetles feed on nectar and pollen, usually by crawling over flower surfaces. There are around 30,000 species of beetles in Australia, with many yet to be formally described.



Butterflies have wings covered in tiny scales. They have clubbed antennae and hold their wings upright when at rest. They are day-flying and have long tongues that they can use to feed on nectar in flowers with deep tubes. Butterflies are usually brightly coloured, with approximately 600 species found in Australia.



Moths also have wings covered in tiny scales and tend to be subtle in colour. They have antennae without clubs and hold their wings flat when at rest. They are generally dusk- and night-flying but there are some exceptions: the grapevine moth is a commonly seen day-flying moth. Moths feed on nectar. Australia has a high diversity of moth species, with up to 22,000 species thought to exist across the continent.

Flower forms



Generalist flowers can be pollinated by many different insects and animals. They are typically saucer shaped with many stamens and have a surface that insects can walk on. *Eucalyptus* flowers and daisy flowers are generalist flowers – they can be pollinated by bees, flies, beetles and butterflies.



Specialist flowers have modifications to their shape and size that only let certain pollinators access the nectar and pollen. These flowers might have deep flower tubes or narrow entry points so that only a select group of pollinators can access them. The advantage of specialisation is that pollination is very targeted and efficient, with accurate pollen placement made possible by co-evolution between flowers and insects. The disadvantage is that if the correct pollinator isn't there, the flowers aren't pollinated. Often, nectar is produced at the base of the flower, forcing pollinators to enter the flower fully and in the process, become covered in pollen.

Pollinator rewards

Nectar is a sugary solution, rich in carbohydrates, vitamins and minerals, produced by flowers and sometimes by glands on leaves or stems (called extra-floral nectaries). Nectar is attractive to insects, and provides an immediate energy source needed for tasks such as hunting pest insects, laying eggs in decomposing organic matter, collecting pollen, or parasitising other insects.

Carbohydrates alone don't support everything needed for health and growth, so insects also need pollen.

Pollen is rich in protein, fats and nutrients. Bees are vegetarian, and need to collect pollen to feed their offspring.

Buzz pollination

Some flowers do not produce any nectar; they specifically target pollencollecting bees, and only offer pollen rewards. To limit pollen loss and ensure effective pollination, some plants produce flowers with specialised, tubular anthers, that only open at the tip. To extract pollen, bees use vibrations to 'buzz' the pollen grains out of the pores of these anthers. Many crops are buzz pollinated, including tomatoes, potatoes, eggplants, capsicum, chillies, tomatillo and cranberries.

European honey bees are unable to buzz pollinate flowers, but several native bees, such as the blue-banded bee, teddy bear bee (*Amegilla* sp.) and carpenter bee (*Xylocopa* sp.) are exceptionally good large buzz pollinators, and have evolved to pollinate native plants such as flax lilies (*Dianella* sp.). Many of our smaller, ground nesting bees utilise vibration to help them excavate their burrows, and they also use that skill to buzz pollen from the anthers of native plants.

Planting buzz-pollinated species will encourage populations of buzz pollinators for successful pollination of food crops and ensure seed set in native plants. Many small ground nesting bees also buzz pollinate native flowers.

Nectar feeding

Grevillea flowers and other tubular flowers are often adapted to be successfully pollinated by birds. Pollen is 'presented' on a floral stigma that extends outside the flower. When birds feed on the nectar, pollen is deposited on their beaks or heads. Bees, also attracted to the sugary nectar, crawl into the side of the flower and feed on the nectar without encountering the pollen-laden stigma. The plant doesn't receive the pollination benefit from the insect, but flowers such Grevillea species can be a very useful source of nectar for insects in the cooler months.



Nurseries

Most of the plants shown in the planting guide will be available at nurseries that have a good stock of native plants. But if your local nursery doesn't stock the plant you're after, ask them to order it in. For a list of nurseries

that stock all the plants shown in the planting guide, plus other useful resources, visit the Wheen Bee Foundation website or scan the QR code.



WheenBeeFoundation.org.au/our-work/powerful-pollinators

Wheen Bee Foundation

Powerful Pollinators Planting Guides are produced by Wheen Bee Foundation. We fund vital strategic research and education initiatives that strengthen bees, improve pollination efficiency, and protect our food security and ecosystem health. Visit the website for more information.

WheenBeeFoundation.org.au

Far left: The spreading flax lily, *Dianella revoluta,* is buzz pollinated.

Left: This European honey bee is 'side-working': feeding on the nectar-rich flowers without coming into contact with the plant's pollen.

Front cover:

 A Lassioglossum bee foraging for pollen on Arthropodium milleflorum.
(Photo: Stan Wawrzyczek)
The Grampians (Gariwerd), viewed from the track to the peak of Mt Stapylton.
(Photo: Stan Wawrzyczek)
European honey bees, Apis mellifera. (Photo: Kirrily Hughes)

This project is supported by: Smart Farms, through funding from the Australian Government's National Landcare Program and La Trobe University.







latrobe.edu.au





PP-VIC30.1 (2023) Wheen Bee Foundation 2023. Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License. To view a copy of this license, visit http://creativecommons.org/licenses/by-nc-nd/4.0/