Powerful pollinators

Encouraging insect pollinators in rural and urban landscapes



Pollinators are an essential component of healthy, biodiverse landscapes and provide critical pollination services to native flora and agriculture production across the country.

This guide provides information on ways to encourage a diverse range of insect pollinators across all properties, and includes a planting calendar to help select plants to support diverse pollinators throughout the year.



The power of pollinators

Pollinators – mostly insects, but also birds and mammals – assist the production of seeds and fruit in many plant species by visiting flowers in search of food (nectar and/or pollen). Whilst foraging they transfer pollen from one flower to another, facilitating fertilisation, which results in fruits and seeds.

Honey bees, native bees and other native insects like hoverflies, wasps and butterflies provide essential pollination services for native plants, pastures, crops, fruits and vegetables.



Native vegetation supports pollinators by providing food and nesting sites. Nearby crops and pastures will benefit from the increased abundance and diversity of pollinators in the landscape.

Pollinators and food security

Without insect pollinators, the quantity and diversity of food grown for humans in contemporary agricultural systems would be severely restricted. Many of the food crops we eat, as well as pasture and fodder crops, benefit from pollination by insects.

Pollinator-dependent crops include avocados, mangos, macadamias and vegetables, as well as many crops grown for seed production, such as legumes and pasture.

The quantity and diversity of insect pollinators are key drivers of production as they influence both crop yields and quality. Under-pollination results in smaller and misshapen fruit or seed that isn't viable.

Grazing enterprises can also suffer from a reduction in the abundance or diversity of pollinators, due to the role these insects play in the persistence of nitrogen-fixing pasture legumes such as clover.

A diverse and healthy community of pollinators generally provides more effective and consistent pollination than relying on any single species. Insect populations are in decline worldwide due to land clearing, intensive or monocultural agriculture, pesticide use, pollution, colony disease, increased urbanisation and climate change. Low pollinator numbers mean not all flowers are pollinated, leading to low fruit or seed set. This in turn reduces fruit and vegetable harvest yields, and decreases food supply.



Under-pollination results in smaller, misshapen fruit such as this strawberry.

Healthy ecosystems

Pollinators are both essential to, and depend upon, healthy ecosystems. A growing human population and increasing demand for food puts pressure on ecosystems, while declining ecosystem function will in turn negatively impact food production.

Insect pollinators are a prime example of this — without healthy ecosystems and the presence of patches of native vegetation to support insect populations, pollination will decline. This will threaten both crop productivity and the persistence of native, pollinator-dependent flowering plants.

Pollinators require habitat that contains year-round food sources, breeding resources and nesting sites. The presence of pollinator habitat adjacent to food crops has been shown to improve food production by enabling a greater variety and number of pollinators to persist year-round, providing pollination services when required.

Turn to the centre of this brochure for a guide to planting for pollinators.

Diapause or diet? Where are the insects?

Many insect pollinators undergo a diapause during colder winter months. Diapause is a period of suspended development during unfavourable environmental conditions, and during this period insect pollinators do not need flowers. Birds and other small mammals will continue to benefit from available pollen and nectar during this time.

If there are low numbers of insect pollinators in your local area, it is important to determine whether this is because of diapause, or because of an inadequate availability of nectar and pollen, creating a 'food desert' where insect pollinators cannot survive.

There are still many unknowns about insect pollinators in Australia. Take part in Australian Pollinator Week or in the annual Australian Pollinator Count to learn more about pollinators in your area – visit:

AustralianPollinatorWeek.org.au and **AustralianPollinatorCount.au**

1

Encouraging pollinators on your property

Create pollination reservoirs

Pollination reservoirs are areas of native plant species that provide floral resources for pollinators. They can be new plantings or existing habitat, such as shelterbelts or remnant vegetation. A high diversity of plant species is essential to provide nectar, pollen and nesting sites throughout the year. Pollination reservoirs need to be close enough to crops to ensure that pollinators can fly easily to them.

Use existing habitat

Protect and improve existing habitat where possible. Roadsides, shelterbelts, dam margins, woodlands, grasslands, rocky areas, river and creek edges can all be important pollinator-attracting areas, bringing valuable pollination services to your property.

Native vegetation stands provide habitat for pollinators, and special attention should be paid to enhance and protect these areas.

Get to know your local flora

Each property and region will have distinct populations of insects, based on the plants and climate. Identifying and understanding the insects in your area will help you develop better plantings.

The plants growing in nearby bush will be well suited to the climate and soils in your region. Local community groups and specialist native nurseries can provide useful information and usually produce local plant species.

Plant trees, shrubs and groundcovers

Planting a variety of species of groundcovers, shrubs and trees on your property will further attract pollinators to your area. Use a combination of direct seed sowing and planting tube stock to establish new vegetation. Initial watering and protection from grazing will improve the success rate of young plants. Wildflowers, including our native pea species, are excellent at attracting a diverse range of native pollinators.

Connectivity counts

Insect pollinators benefit from greater connectivity of habitat in a landscape, which allows them to forage over a wider radius and increase in numbers in a local area. Encourage neighbours and other landholders to plant for pollinators and create connections across your landscape.

Utilise ecotones

Ecotones are the margins between two different habitats. Ecotones often contain a more diverse mixture of pollinator species because they are inhabited by pollinators from both habitats. Protect and utilise ecotones such as the transition zones between woodland and grassland, or heath and shrubland, to create highly diverse floral and insect communities.

Amplify the flower signal

Plants have evolved large flowers or clusters of smaller flowers which attract more pollinator visits. Large, colourful and diverse plantings attract more pollinators. Ideally, plant in groups that contain different vegetation layers — combine a species-rich mixture of wildflowers, groundcovers, herbs, lilies, rushes, climbers, shrubs and trees.

Plant for the future

When establishing pollinator habitat, consider including species that are indigenous to your area and can tolerate increasingly warmer and drier (or wetter) conditions, to improve resilience to climate change. Rehabilitate weedy areas into managed pollination reservoirs by introducing lots of flowering plant diversity.

Be careful not to plant invasive or listed weeds, and look for suitable replacements.

Double the crop value

Plants that are pollinator-attracting may be commercially viable crop species in their own right and can be used to diversify farm production. Bush foods such as Native Ginger, Wild Tamarind, Lemon Aspen and many more are in high demand for use in fresh and manufactured products. Native plant seed is also needed for revegetation projects. Farmers can also support beekeepers by hosting beehives to increase pollinator numbers on a farm.

Reduce chemical use

Insecticides, fungicides and herbicides all affect the health of bees, bee colonies and native pollinators.
Herbicides can impact pollinators by reducing the availability and diversity of flora and removing vegetation that helps support insect life. Some herbicides can also harm the beneficial microbes in the insect gut. In many circumstances, beneficial insects will, in healthy numbers, help control pest insects, ultimately reducing the need for insecticide use.

When chemical pest control is unavoidable, select products that are least harmful for pollinators and apply insecticides in the evening or at night when pollinators are not active.

Always use according to directions, especially for withholding periods, and notify beekeepers a few days before spraying chemicals so beehives can be safely relocated away from harm.

Be a citizen scientist and do some detective work to discover local pollinators on your property. Visit inaturalist.ala.org.au to be involved.

Safeguard the bees? The best way to 'save the bees' and protect our pollinators is to create an abundance of diverse habitat — from the ground up! There is much interest in keeping a beehive to promote pollinators, but there are serious legal and biosecurity responsibilities that must be considered, and that the introduction of a beehive does not displace existing native pollinators and insects. Be a friend of pollinators and say it with flowers!

A guide to planting for pollinators for the Atherton Tablelands region, Queensland



Healthy populations of insect pollinators are important for crop yields, orchard production and thriving native vegetation.

This planting guide will help you choose plant species to attract and keep pollinators on your property throughout the year.

The Atherton Tablelands encompasses both the Wet Tropics Bioregion and the Einasleigh Uplands Bioregion. This is a diverse area spreading west from the coastal escarpment of Cairns, Queensland. The mountainous region reaches peaks above 1600 m, the highest in Queensland. The Atherton Tablelands has recorded some of the highest annual rainfall in Australia. Soils in the region, are often derived from basalt and are highly fertile and well-drained. The elevation, climate and soils support prolific native biodiversity, including tropical rainforests and savannahs. This area supports a wide range of agricultural farms, supplying both the domestic and international markets.

The plants listed in this Guide will help supply rewards to pollinators, with an emphasis on species that are indigenous and suited to local climates.

The eucalypt species in the chart have been selected as high quality honey production species. Most eucalypts do not flower every year, so choosing diverse species will help create continuously flowering habitat.



The pollinator plant list

To create pollinator-attracting plantings, use the Guide to choose a selection of plants with a variety of flower colours, different growth habits and a range of flowering seasons.

For each species, the planting Guide lists:

- life-form/'habit' (climber, herb, shrub or tree) and height (m).
- the vegetation type in which they naturally occur
- flower colour and flowering season
- growth requirements (sun/shade, moist/dry)
- insect groups that may visit each plant and the floral reward (pollen and/or nectar).

The coloured bars indicate the flowering months for each species. Darker shading denotes the peak flowering period, with a lighter shading for non-peak flowering months. Flowering dates may differ between regions and seasons, particularly for non-peak times, if your local climate is consistently warmer or cooler than average, with earlier or later flowering.

Sourcing plants

Most of the plant species listed are available from retail or wholesale nurseries or native plant growers, and local environment groups. If you can't source these plants at your local garden centre, or indigenous nursery, ask them to contact the local nursery suppliers and plant growers listed online. See the reverse of the Guide for details.

WheenBeeFoundation.org.au

Lifeform	C	Scientific name	Family	Vogatation type	Height	Flower colour Flowering		Annat	Sail maiatura	Pollinato	r reward		Visitation by pollinator					
	Common name	Scientific name	Family	Vegetation type	Height	Flower colour	Jan Feb Mar Apr May Jun Jul Aug Sep	Oct Nov Dec Aspect	Soil moisture	Pollen	Nectar	Native bees	Honey bees	Hoverflies	Wasps E	Butterflies	Moths Be	eetles
Crop plants																		
orbs / Herbs	Native Banana	Musa banksii	Musaceae	Lowland, Upland Rainforest	6 m	Cream-Yellow		Full sun to light shad		•	•		•					•
orbs / Herbs ee	Tea Plant	Camellia sinensis	Theaceae	Cultivated	1 m	Cream, Yellow		Full sun to light shad		•	•	•	•	•	•			•
	Macadamia Mango 'Kensington Pride'	Macadamia integrifolia	Proteaceae Anacardiaceae	Sub-Tropical Rainforest Cultivated	5–12 m < 24 m	Cream Cream-Yellow		Full sun to light shace	de Moist, well-drained Well-drained	•	•			•	•			•
nes & Climbers	Passionfruit	Mangifera indica Passiflora edulis	Passifloraceae	Cultivated	Climber	Purple		Full sun	Dry to moist					•				
Indigenous plants	Passioninuii	Passiliora eaulis	Passilloraceae	Cullivaled	Cilifiber	Purple		Full Sull	Dry to moisi									
orbs / Herbs	Native Ginger	Alpinia caerulea	Zingiberaceae	Sub-Tropical Rainforest	2-2.5 m	White		Sun to full shade	Moist, well-drained		•							
orbs / Herbs	Blue Flax Lily	Dianella atraxis	Asphodelaceae	Sclerophyll Forest	0.8 m	Purple		Sun to semi-shade	,	*		•						
orbs / Herbs	Blue Flax-Lily	Dianella longifolia	Asphodelaceae	Sclerophyll Forest	0.8 m	Purple		Sun to semi-shade		*		•						
orbs / Herbs	Settlers' Twine	Gymnostachys anceps	Araceae	Rain & Wet Sclerophyll Forest, Coastal	< 3 m	Cream, Green, Yellow		Full shade	Moist	•	•	•	•	•	•			•
orbs / Herbs	Small St. John's Wort	Hypericum gramineum	Hypericaceae	Open Forest, Grassland	0.1-0.4 m	Yellow		Sun to semi-shade		•		•	•	•	•			
orbs / Herbs	Birdsville Indigo	Indigofera linnaei	Fabaceae	Wood & Shrubland, Wooded Grassland	0.15-0.5 m	Red		Sun to semi-shade	Dry, well-drained	•	•	•	•			•		•
orbs / Herbs	Pastel Flower	Pseuderanthemum variabile	Acanthaceae	Rainforest, Wet Scleorphyll Forest	0.15-0.3 m	White, Mauve		Sun to light shade	Moist, well-drained	•	•	•				•		•
orbs / Herbs	Arrowroot	Tacca leontopetaloides	Dioscoreaceae	Rainforest, Monsoon Forest	<1m	Purple, Green, Brown		Semi-shade to full sha	de Moist, well-drained	•								
orbs / Herbs	Trephosia	Tephrosia filipes	Fabaceae	Sclerophyll Forest	< 2 m	Purple		Full sun	Dry, well-drained	•	•	•	•			•		
rass / Sedge / Rush	Many-Flowered Mat-Rush	Lomandra multiflora	Asparagaceae	Forest, Woodland	0.2-0.9 m	Cream		Sun to semi-shade	Variable	•	•	•	•		•			•
rass / Sedge / Rush	Awned Panick Grass	Oplismenus compositus	Poaceae	Forest	0.3-0.8 m	Pink		Semi-shade to shad	de Moist	•								•
lies & Irises	Stream Lily	Crinum pedunculatum	Amaryllidaceae	Forest, Wetland	0.3-2 m	White		Sun to semi-shade	Moist	•	•	•		•			•	•
nrub (Small)	Wattle	Acacia umbellata	Fabaceae	Eucalypt Woodland, Rocky Slopes & Ridges	< 3 m	Golden		Full sun	Dry, well-drained	•	•	•	•	•	•	•		•
rub (Small)	Great Woolly Malayan Lilac	Callicarpa candicans	Lamiaceae	Rainforest, Vine Thickets, Monsoon Forest	2 m	Purple		Sun to semi-shade	Dry, well-drained	•		•	•	•	•	•	•	•
nrub (Small)	Gardenia White Star	Gardenia psidioides	Rubiaceae	Eucalypt & Monsoon Forest, Vine Thickets	1 m	White		Sun to semi-shade	Well-drained	•	•					•	•	
nrub (Small)	Tinaroo Bottlebrush	Melaleuca recurva	Myrtaceae	Sclerophyll & Open Forest	< 7 m	Red		Sun to semi-shade	Variable	•	•	•	•	•				
nrub (Small)	Blue Tongue	Melastoma malabathricum	Melastomataceae	Rain, Monsoon & Wet Open Forest, Heath	1–3 m	Pink		Full sun to full shac	le Moist, well-drained	•*		•	•					
rub (Small)	Phyllanthus	Phyllanthus lamprophyllus	Phyllanthaceae	Open & Monsoon Forest	1–2 m	Cream		Sun to semi-shade	Dry, well-drained	•	•						•	
rub (Large)	Heathlands Wattle	Acacia simsii	Fabaceae	Eucalypt Open Forest, Woodland	2-4 m	Golden		Full sun	Dry, well-drained	•		•	•	•	•	•		•
rub (Large)	White Ball Acacia	Acaciella angustissima	Fabaceae	Savannahs, Woodlands	2-7 m	Cream		Sun to semi-shade	Dry-moist, well-drained	•	•	•	•	•	•	•	•	•
rub (Large)	Mock Orange	Bursaria tenuifolia	Pittosporaceae	Riparian Forest and Scrubland	3-9 m	White & Green		Light to moderate sha	de Moist-dry, well-drained	•	•	•	•	•	•	•	•	•
rub (Large)	Apricot Myrtle	Pilidiostigma tropicum	Myrtaceae	Dry Rainforest	1–5 m	Cream-Yellow	<u> </u>	Sun to semi-shade	Moist, well-drained	•	•	•	•	•	•	•	•	•
rub (Large)	Wild Yellow Jasmine	Pittosporum revolutum	Pittosporaceae	Rain, Wet Sclerophyll & Open Forest	< 3 m	Yellow		Sun to semi-shade	Moist, well-drained	•	•	•	•	•	•	•	•	•
rub (Large)	Beach Naupaka	Scaevola taccada	Goodeniaceae	Sandy Beaches, Rocky Outcrops	< 4 m	Cream		Sun to light shade	Well-drained	•	•	•	•			•		
rub (Large)	Golden Penda	Xanthostemon chrysanthus	,	Coastal Rainforest	10-15 m	Golden		Full sun	Moist, well-drained	•	•	•	•	•	•	•		•
ree (Small)	Club-Leaf Wattle	Acacia hemignosta	Fabaceae	Open Woodland	3–10 m	Golden		Full sun	Dry, well-drained			•	•	•	•	•		•
ree (Small)	Lemon Aspen	Acronychia acidula	Rutaceae	Rainforest	12–15 m	Cream		Full sun	Moist, well-drained		•	•	•	•	•	•	•	•
ree (Small)	Native Gardenia	Atractocarpus fitzalanii	Rubiaceae	Rainforest	< 6 m	Cream		Sun to half shade	Moist, well-drained		•	•	•		•			•
ree (Small)	Jigal Tree	Lysiphyllum cunninghamii	Fabaceae	Open Forest	< 12 m	Red		Full sun	Dry, well drained		•	•				•		
ree (Small)	Weeping Bottlebrush	Callistemon viminalis	Myrtaceae	Riparian forest	6-9 m	Red		Sun to semi-shade			•	•	•	•				
ree (Small)	Red Bloodwood	Corymbia erythrophloia	Myrtaceae	Open Eucalyptus Woodland	< 8 m	Cream		Full sun	Dry, well-drained		•	•	•	-	•	•	•	•
ree (Small)	Tuckeroo	Cupaniopsis anacardioides	· '	Littoral Rainforest, Scrub	< 12 m	Cream, Green, Pink		Full sun	Moderate, well-drained		•	•	•	•	•	•		•
ree (Small)	Broad-Leaved Ballart	Exocarpos latifolius	Santalaceae	Variable, Open habitats	2–10 m	Green		Sun to semi-shade	,,			•	•	•	•	•		•
ree (Small)	Python Tree	Gossia bidwillii	Myrtaceae	Rainforest	18 m	White		Semi to full shade	Moist, well-drained		•	•	•		•	•		•
ree (Small)	Fern-Leaf Grevillea	Grevillea pteridifolia	Proteaceae	Open Forest Rainforest	4-8 m	Golden		Full sun	Dry, well-drained Well-drained		•	•	•		•			•
ee (Small)	Canary Beech	Huberantha nitidissima	Annonaceae	Rainforest	< 10 m	Cream-Yellow Cream-Yellow		Sun to semi-shade Sun to semi-shade		•	•		•		•			•
ee (Small) ee (Small)	Kamala Myrtle	Macaranga tanarius Melaleuca borealis	Euphorbiaceae Myrtaceae	Open Forest, Woodland	< 10 m	Cream Cream		Sun to semi-shade	· ·		•			•	•			
ee (Small)	Hard Pink Alder	Ostrearia australiana	Hamamelidaceae	'	1–6 m			Semi-shade	Moist, well-drained		•							•
	White Siris		Fabaceae	Open, Rain & Monsoon Forest	< 25 m			Sun to light shade	Dry, well-drained									
ee (Large) ee (Large)	Alectryon	Albizia procera Alectryon connatus	Sapindaceae	Variable Forests	10-20 m			Sun to half shade	Variable, well-drained		•	•		•				•
ee (Large)	Atherton Oak	Athertonia diversifolia	Proteaceae	Coastal Ranges	5–10 m	Cream Cream		Sun to light shade	Well-drained					•	•			
ee (Large)	Illawarra Flame Tree	Brachychiton acerifolius	Malvaceae	Subtropical Rainforest	< 35 m	Red		Sun to semi-shade		•	•	•	•	•		•		•
ee (Large)	Cinnamon Laurel	Cryptocarya densiflora	Lauraceae	Mountain Rainforest	7–20 m	Cream-Green		Moderate to full shad		•	•	•	•	•	•			•
ee (Large)	Wild Tamarind	Diploglottis diphyllostegia	Sapindaceae	Rainforest	< 30 m	Cream		Semi to full shade	Moist, well-drained		•	•	•	•	•	•		•
ee (Large)	Woollybutt	Eucalyptus miniata	Myrtaceae	Dry Sclerophyll Forest	10-20 m	Orange, Yellow, Red		Full sun	Dry, well-drained		•	•	•	•	•	•	-	•
ee (Large)	Forest Red Gum	Eucalyptus tereticornis	Myrtaceae	Wet & Dry Sclerophyll Forest, Woodland		Cream		Sun to semi-shade	,,,		•	•	•	•	•	•	•	•
ee (Large)	Small-Leaved Fig	Ficus obliqua	Moraceae	Rainforest, Coastal	< 30 m	Inconspicuous		Full sun	Variable	•					•			
ee (Large)	Teak	Flindersia australis	Rutaceae	Dry & Subtropical Rainforest	< 40 m	White		Sun to semi-shade		•	•	•	•	•	•	•		•
e (Large)	Cascading Bean	Maniltoa lenticellata	Fabaceae	Tropical Rainforest	< 22 m	Cream		Sun to semi-shade			•	•	•	•	•	•	•	•
e (Large)	Pink-Flowered Doughwood	Melicope elleryana	Rutaceae	Subtropical Rainforest, Swamp Forest	< 25 m	Pink		Sun to semi-shade			•	•	•	•	•	•		•
e (Large)	Brush Cherry	Syzygium australe	Myrtaceae	Forest	< 35 m	White		Sun to semi-shade	-	•	•	•	•	•	•	•	•	•
e (Large)	Kuranda Satinash	Syzygium kuranda	Myrtaceae	Rainforest	< 35 m	White		Full sun	Moist, well-drained	•	•	•	•	•	•	•	•	•
e (Large)	Damson Plum	Terminalia sericocarpa	Combretaceae	Dry & Monsoon Rainforest, Rainforest	10-40 m	White		Sun to semi-shade	,		•	•	•	•	•			•
e (Large)	Red Cedar	Toona ciliata	Meliaceae	Wet & Dry Sclerophyll Forest, Rainforest		Pink		Sun to light shade	Moist, well-drained	_	•	•	•	•	•			•
es & Climbers	Scrambling Lily	Geitonoplesium cymosum	Asphodelaceae	Sclerophyll Forest, Woodland	Climber	White		Semi to full shade	Dry, well-drained		•	•	•	•				•
nes & Climbers	Climbing Guinea Flower	Hibbertia scandens	Dilleniaceae	Rainforest	Climber	Yellow		Sun to half shade	Dry, well-drained			•	•	•	•			
es & Climbers	Jasmine	Jasminum simplicifolium	Oleaceae	Drier Rainforest, Monsoon Forest	Climber	White		Sun to semi-shade	Variable	•	•	•	•	•	•	•	•	•
		,		Farest Manadlerad	Climber			Com to comi alondo	Moist to dry	•			•	•				•
nes & Climbers	Wonga Wonga Vine	Pandorea pandorana	Bignoniaceae	Forest, Woodland	Cilifiber	writte-rurple		Sun to semi-shade	i Moisi io di y					_				_

Know your pollinators



European honey bees have two pairs of wings and long, segmented antennae. They are daytime-flying and feed on nectar and pollen. They are generalist pollinators and provide the bulk of pollination services for horticulture and crop plants. Honey bees and native bees are both essential to functioning ecosystems and food security in Australia.

Honey bees have become an important part of the Australian landscape. Honey bees live as colonies, and have a long history of coexistence with humans, including in domestic gardens.



Australian native bees comprise more than 2000 species, which provide essential pollination services. Native bees are generally solitary and live in nests in the ground or in hollow stems, old borer holes and other cracks and crevices, and some have evolved to pollinate particular native flowers through 'buzz pollination'. Although many Australian native bees are generalist foragers, some species have co-evolved with native plants and adapted to be the most effective pollinators of their flowers. Many native plant species, such as Dianella and Grevillea require specially adapted insects to access their nectar and enable the transfer of pollen to the stigma. Most native bees are solitary, but some species found in northern Australia (Tetragonula sp. and Austroplebeia sp.) are social bees and are used for commercial pollination of crops like macadamia nuts.



Fly species number up to 30,000 in Australia, and can be identified by having only one pair of flight wings. A second set of wings are modified into club-shaped paddles that allow flies to hover and stabilise their flight. Unlike bees and wasps, many flies (Brachycera) have very small, clubbed antennae at the front of their head. Flies, including blowflies, are often attracted to flowers that smell like carrion. Some flower-flies, have hairy bodies that easily collect pollen while they are feeding. Flies provide a range of services in the garden, including pollination, decomposition and predation.



Hoverflies are a type of fly, distinguishable by their large eyes, short antennae, bright black and yellow abdomen and their hovering flight behaviour. Adult hoverflies are nectar and pollen feeders. Hoverfly larvae feed on pests such as aphids, thrips and leafhoppers and are excellent biocontrol agents.



Beetles have hard outer wings that form their distinctive beetle shape. Their outer wings form a T-shape where they join at the top, unlike bugs where the outer wings make an X- or Y-shape. Some beetles feed on nectar and pollen, usually by crawling over flower surfaces. There are around 30,000 species of beetles in Australia, with many yet to be formally described.



Butterflies have wings covered in tiny scales. They have clubbed antennae and hold their wings upright when at rest. They are day-flying and have long tongues that they can use to feed on nectar in flowers with deep tubes. Butterflies are usually brightly coloured, with approximately 600 species found in Australia.



Moths also have wings covered in tiny scales and tend to be subtle in colour. They have antennae without clubs and hold their wings flat when at rest. They are generally dusk- and night-flying but there are some exceptions: the grapevine moth is a commonly seen day-flying moth. Moths feed on nectar. Australia has a high diversity of moth species, with up to 22,000 species thought to exist across the continent.

Flower forms



Generalist flowers can be pollinated by many different insects and animals. They are typically saucer shaped with many stamens and have a surface that insects can walk on. Eucalyptus flowers and daisy flowers are generalist flowers — they can be pollinated by bees, flies, beetles and butterflies.



Specialist flowers have modifications to their shape and size that only let certain pollinators access the nectar and pollen. These flowers might have deep flower tubes or narrow entry points so that only a select group of pollinators can access them. The advantage of specialisation is that pollination is very targeted and efficient, with accurate pollen placement made possible by co-evolution between flowers and insects. The disadvantage is that if the correct pollinator isn't there, the flowers aren't pollinated. Often, nectar is produced at the base of the flower, forcing pollinators to enter the flower fully and in the process, become covered in pollen.

Pollinator rewards

Nectar is a sugary solution, rich in carbohydrates, vitamins and minerals, produced by flowers and sometimes by glands on leaves or stems (called extra-floral nectaries). Nectar is attractive to insects, and provides an immediate energy source needed for tasks such as hunting pest insects, laying eggs in decomposing organic matter, collecting pollen, or parasitising other insects.

Carbohydrates alone don't support everything needed for health and growth, so insects also need pollen.

Pollen is rich in protein, fats and nutrients. Bees are vegetarian, and need to collect pollen to feed their offspring.

Buzz pollination

Some flowers do not produce any nectar; they specifically target pollen-collecting bees, and only offer pollen rewards. To limit pollen loss and ensure effective pollination, some plants produce flowers with specialised, tubular anthers, that only open at the tip. To extract pollen, bees use vibrations to 'buzz' the pollen grains out of the pores of these anthers. Many crops are buzz pollinated, including tomatoes, potatoes, eggplants, capsicum, chillies, tomatillo and cranberries.

European honey bees are unable to buzz pollinate flowers, but several native bees, such as the blue-banded bee, teddy bear bee (Amegilla sp.) and carpenter bee (Xylocopa sp.) are exceptionally good large buzz pollinators, and have evolved to pollinate native plants such as flax lilies (Dianella sp.). Many of our smaller, ground nesting bees utilise vibration to help them excavate their burrows, and they also

use that skill to buzz pollen from the anthers of native plants.

Planting buzz-pollinated species will encourage populations of buzz pollinators for successful pollination of food crops and ensure seed set in native plants. Many small ground nesting bees also buzz pollinate native flowers.

Nectar feeding

Grevillea flowers and other tubular flowers are often adapted to be successfully pollinated by birds. Pollen is 'presented' on a floral stigma that extends outside the flower. When birds feed on the nectar, pollen is deposited on their beaks or heads. Bees, also attracted to the sugary nectar, crawl into the side of the flower and feed on the nectar without encountering the pollen-laden stigma. The plant doesn't receive the pollination benefit from the insect, but flowers such Grevillea species can be a very useful source of nectar for insects in the cooler months.





Nurseries

Most of the plants shown in the planting guide will be available at nurseries that have a good stock of native plants. But if your local nursery doesn't stock the plant you're after, ask them to order it in. For a list of nurseries that stock all the plants shown in the planting guide, plus other useful resources, visit the Wheen Bee Foundation website or scan the QR code.

WheenBeeFoundation.org.au/our-work/powerful-pollinators

Wheen Bee Foundation

Powerful Pollinators Planting Guides are produced by Wheen Bee Foundation. We fund vital strategic research and education initiatives that strengthen bees, improve pollination efficiency, and protect our food security and ecosystem health. Visit the website for more information.

WheenBeeFoundation.org.au

Far left: The spreading flax lily, Dianella revoluta, is buzz pollinated.

Left: This European honey bee is 'side-working': feeding on the nectar-rich flowers without coming into contact with the plant's pollen.

Front cover:

- Hyleaus species on Melastoma
 malabathricum L. subsp. malabathricum.

 (Photo: Jaura Lagreti)
- 2. Mount Bartle Frere. (Photo: Laura Lopresti)
- 3. European honey bees,

 Apis mellifera. (Photo: Kirrily Hughes)

This project is supported by: Smart Farms, through funding from the Australian Government's National Landcare Program and James Cook University.







jcu.edu.au



