Powerful pollinators

Encouraging insect pollinators in urban environments & gardens



Pollinators are an essential component of our cities and suburbs. Enhancing pollinator resources to support a diverse range of beneficial insects is important for flourishing gardens, sustainable backyards, healthy green spaces, and thriving ecosystems. The Powerful Pollinators Guide provides an introduction to encouraging insect pollinators in the south-eastern suburbs and surrounds of urban Melbourne.



The power of pollinators

Pollinators – mostly insects, but also birds and mammals – assist the production of seeds and fruit in many plant species by visiting flowers in search of food (nectar and/or pollen). Whilst foraging they transfer pollen from one flower to another, facilitating fertilisation, which results in fruits and seeds.

Honey bees, native bees and other native insects like hoverflies, wasps and butterflies provide essential pollination services for native plants, garden flowers, fruits and vegetables.

Pollinators and food security

Without insect pollinators, the quantity and diversity of food and plants grown in backyard gardens would be severely restricted. Many of the foods we eat, from gardens and farms, benefit from pollination.

Pollinator-dependent foods include citrus, apples, stone-fruits, zucchini, pumpkins, strawberries and tomatoes, as well as plants grown for seed such as sunflowers, coriander and parsley.

The quantity and diversity of insect pollinators are key drivers of production as they influence both food yields and quality. Under-pollination results in smaller and misshapen fruit or seed that isn't viable.

A diverse and healthy community of pollinators generally provides more effective and consistent pollination than relying on any single species.

Pollinators are essential to, and dependent upon, healthy ecosystems. A growing human population and increasing demand for food puts pressure on ecosystems, with potential negative impacts on biodiversity, the environment and food production.



Increasing the abundance and diversity of native plants in urban landscapes supports pollinators by providing a range of food sources and nesting sites.

Insect populations are in decline worldwide due to land clearing, intensive or monocultural agriculture, pesticide use, pollution, colony disease, increased urbanisation and climate change. Low pollinator numbers mean not all flowers are pollinated, leading to low fruit or seed set. This in turn reduces fruit and vegetable harvest yields, and decreases food supply.



Under-pollination results in smaller, misshapen fruit such as this strawberry.

Backyard biodiversity

Insect pollinators are a prime example of the importance of healthy ecosystems in urban gardens, parks and reserves. Insects are the 'canaries in the coal mine' of our urban and rural environments. Without our 'littlest creatures', we lack pollinators, natural beneficial pest control services, and critical food source for other insects, birds, amphibians, reptiles and mammals.

The presence of connected and widespread pollinator habitat is critical to support insect populations if we are to maintain sustainable cities and productive, healthy gardens and urban farms for food security and biodiversity.

Pollinators require habitat that contains year-round food sources, breeding resources and nesting sites. The presence of pollinator habitat adjacent to food crops has been shown to improve food production by enabling a greater variety and number of pollinators to persist year-round, providing pollination services when required.

Turn to the centre of this brochure for a guide to planting for pollinators.

Diapause or diet? Where are the insects?

Many insect pollinators undergo a diapause during colder winter months. Diapause is a period of suspended development during unfavourable environmental conditions, and during this period insect pollinators do not need flowers. Birds and other small mammals will continue to benefit from available pollen and nectar during this time.

If there are low numbers of insect pollinators in your local area, it is important to determine whether this is because of diapause, or because of an inadequate availability of nectar and pollen creating a 'food desert' where insect pollinators cannot survive.

There are still many unknowns about insect pollinators in Australia. Take part in Australian Pollinator Week or in the annual Australian Pollinator Count to learn more about pollinators in your area – visit: **AustralianPollinatorWeek.org.au** and **AustralianPollinatorCount.au**

Encouraging pollinators in your garden

Create pollination reservoirs

Pollination reservoirs are areas that provide floral resources for pollinators. They can be gardens, new plantings or existing habitat such as established trees, or even local bushland, parks or reserves. A high diversity of plant species is essential to provide nectar, pollen and nesting sites throughout the year. Pollination reservoirs need to be close enough to where pollinators live to ensure that they can fly easily to them.

Improve on what you have

Enhance and improve existing pollinator habitat where possible. Gardens that already contain elements such as large trees, rockeries, water sources, bare soil and organic matter, and a variety of flowering plants are a valuable resource for beneficial insects and pollinators.

Nature-strips and verges, ponds, vegetable gardens, orchards, bushland, grassland, and river and creek edges can all be important pollinator-attracting areas. Protect and enhance any native pollinator plants in your garden or on your property for the future.

Plant trees, shrubs and groundcovers

Planting a variety of species of groundcovers, shrubs and trees in your garden will further attract pollinators to your patch. Initial watering and protection will improve the success rate of young plants. Wildflowers, including our native pea species, are excellent at attracting a diverse range of native pollinators.

Be a citizen scientist and do some detective work to discover local pollinators in your patch. Visit **inaturalist.ala.org.au** to be involved.

Construct insect real estate

Insect hotels, which are both functional and attractive, are a great way to add to habitat and nesting places for pollinators and insects in your backyard or garden. The hotels are easily moved to be close to flowering plants and those needing pollination, especially if you have a new garden that is still growing. Include lots of different sized holes, cracks and crevices to provide homes for various solitary insect pollinators.

Plant for the future

When establishing pollinator habitat, consider including species that are indigenous to your area but can tolerate increasingly drier and warmer conditions, to create resilient habitat for climate change. Rehabilitate weedy areas into managed pollination reservoirs by introducing lots of flowering plant diversity. Be careful not to plant invasive or listed weeds, and look for suitable replacements.

Amplify the flower signal

Plants have evolved large flowers or clusters of smaller flowers because they attract more pollinator visits. Large, colourful and diverse plantings attract more pollinators. Ideally, plant in groups that contain different vegetation layers – combine a species-rich mixture of wildflowers, ground-covers, herbs, lilies, rushes, climbers, shrubs and trees.

Connectivity counts

Insect pollinators benefit from greater connectivity of habitat in a landscape, which allows them to forage over a wider radius and increase in numbers in a local area. Encourage friends and neighbours to plant for pollinators and create connections in your community.

Get to know your local flora

Your local government area has distinct populations of insects, depending on the local flora and environment. Knowing your local insect species will help you develop better plantings.

The plants growing in nearby nature reserves or bushland will be suited to your climate and soils. Local environment groups and specialist native nurseries can provide information about local plants.

Grow a bumper crop

Pollinator-attracting plants include many fruits and vegetables grown in backyards, community and market gardens, and orchards. Pollinators ensure good yields of crops such as apples, beans, avocado, and almonds, and bush foods such as lilly-pilly and yam daisy.

Reduce chemical use

Insecticides, fungicides and herbicides all affect bee, colony and wild pollinator health. Herbicides can impact pollinators by reducing the availability and diversity of flora and removing vegetation that helps support insect life. Some herbicides can also harm the beneficial microbes in the insect gut. Insecticides are an obvious threat to pollinators, yet many beneficial insects will, in healthy numbers, help control pest insects, ultimately reducing the need for insecticide use.

Many crops are dependent on pollination by bees. When chemical pest control is unavoidable, select products that are least harmful for pollinators and apply insecticides in the evening or at night when pollinators are not active. Always use according to directions, especially for withholding periods, and notify beekeepers a few days before spraying chemicals so beehives can be safely relocated away from harm.

Safeguard the bees? The best way to 'save the bees' and protect our pollinators is to create an abundance of diverse habitat — from the ground up! There is much interest in keeping a bee hive to promote pollinators, but there are serious legal and biosecurity responsibilities that must be considered, and that the introduction of a bee hive does not displace existing native pollinators and insects. Be a friend of pollinators and say it with flowers!

A guide to planting for pollinators for Metropolitan & Greater Melbourne South East



Healthy populations of insect pollinators are important for sustainable gardens, productive orchards and vegetable crops, and healthy native vegetation.

This Guide will help you select plant species to attract and sustain pollinators • life-form/'habit' (climber, herb, in your garden throughout the year.

The south-eastern Melbourne metropolitan region is defined by mostly alluvial geology, with hilly and undulating topography. The climate is characterised by cool to mild and wet winters, and dry, often hot summers. The vegetation communities that grow naturally include areas of wet forest, woodlands and heathlands, as well as coastal scrub, riparian (streamside) communities, and pockets of coolclimate rainforest.

The plants listed in this Guide have been selected for their resilience and capacity to supply rewards to pollinators, with an differ between regions and seasons, emphasis on species that are indigenous particularly for non-peak times, if your and suited to local climates.

The eucalypt species in this Guide are mostly large trees, but have been included for their value as good nectar producing species. Most eucalypts do not flower every year, so choosing diverse species will help create continuously flowering habitat.



The pollinator plant list

To create pollinator-attracting plantings, use the Guide to choose a selection of plants with a variety of flower colours, different growth habits and a range of flowering seasons.

For each species, the planting Guide lists:

- shrub or tree) and height (m).
- the vegetation type in which they naturally occur
- flower colour and flowering season
- growth requirements (sun/shade, moist/dry)
- insect groups that may visit each plant and the floral reward (pollen and/or nectar).

The coloured bars indicate the flowering months for each species. Darker shading denotes the peak flowering period, with a lighter shading for non-peak flowering months. Flowering dates may local climate is consistently warmer or cooler than average, with earlier or later flowering.

Sourcing plants

Most of the plant species listed are available from retail or wholesale nurseries or native plant growers, and local environment groups. If you can't source these plants at your local garden centre, or indigenous nursery, ask them to contact the local nursery suppliers and plant growers listed online. See the reverse of the Guide for details.

Product of protocols Product	1:6-6	C ommon a a a		Family	Venetation ture	Uninht	Flowerselow		Flowering		Assest	Coil moisture	Pollinato	or reward			Visitatio	n by polline	ator			
Subsect Main and and any and any and any	Lifeform	Common name	Scientific name	Family	Vegetation type	Height		Jc	an Feb Mar Apr May Jun Jul	Aug Sep Oct Nov Dec		Soil moisture	Pollen	Nectar No	ative bees Ho	oney bees	Hoverflies	Wasps Bu	utterflies	Moths	Beetles	Flies
window window window <td></td> <td></td> <td></td> <td></td> <td>,</td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td>· · · · ·</td> <td></td> <td>•</td> <td>•</td> <td>•</td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td>					,							· · · · ·		•	•	•						
Control Standardow Standardow <tt>Standardow</tt> <tt>Standardow<</tt>		0	•		1 1							,	•	•	•	•		•	•		•	
See 1990 See												/		•	•				•	•	•	
Dir Dir <ttt>Dir <tt>Dir</tt> <tt>Dir<td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td>•</td><td>•</td><td>•</td><td>•</td><td></td><td></td><td>•</td><td></td><td>•</td><td></td></tt></ttt>													•	•	•	•			•		•	
Distant Picture					, ,							/	-	-	-						•	
Index Internation Series Seri	Wildflowers	Blue Pincushion			Grassland, Herbfield, Woodland	0.3 m	Blue				Sun	Dry	•	•								
Shipped	Wildflowers	Clustered Everlasting	Chrysocephalum semipapposum	n Asteraceae	Woodland, Heathland	< 0.6 m	Yellow	•			Sun	Moist to dry	•	•	•				•		•	•
		ě	I		, ,			•				/	•	•	•	•	•	•	•		•	•
Subset Subset Subset Subset Subset <td></td> <td>,</td> <td>,</td> <td></td> <td>· · ·</td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td>,</td> <td>-</td> <td>•</td> <td>•</td> <td>•</td> <td></td> <td>•</td> <td>•</td> <td></td> <td>•</td> <td></td>		,	,		· · ·							,	-	•	•	•		•	•		•	
			5		,							· · · ·	-	-	•	•		•	•		•	•
			•		, ,							,		-	•				•		•	
Bits Mar. Show		,		1 0									•*	-	•							
Image Image <t< td=""><td>Lilies & Irises</td><td>Bulbine Lily</td><td>Bulbine bulbosa</td><td></td><td>Grassland</td><td>0.3 m</td><td>Yellow</td><td>•</td><td></td><td></td><td>Sun to semi-shade</td><td>Moist to dry</td><td>•</td><td>•</td><td>•</td><td>•</td><td>•</td><td></td><td>•</td><td>•</td><td>•</td><td>•</td></t<>	Lilies & Irises	Bulbine Lily	Bulbine bulbosa		Grassland	0.3 m	Yellow	•			Sun to semi-shade	Moist to dry	•	•	•	•	•		•	•	•	•
Skale Ale Mark Skale A	Lilies & Irises	Black Anther Flax Lily	Dianella revoluta	Asphodelaceae	Woodland, Heathland	1 m	Indigo				Sun to semi-shade	Moist to dry	•*		•							
Schole State Schole State<		,	Dianella tasmanica	Asphodelaceae	Woodland, Forest		Indigo				Sun to semi-shade	Moist to dry	•*		•							
Biolog Biolog<		•			,									•	•				•	•		
Subject Subject <t< td=""><td>0</td><td></td><td></td><td></td><td>, ,</td><td></td><td></td><td></td><td></td><td></td><td></td><td>/</td><td>-</td><td>-</td><td>•</td><td>•</td><td></td><td>-</td><td>•</td><td></td><td>•</td><td>•</td></t<>	0				, ,							/	-	-	•	•		-	•		•	•
Name Normal Markel		1 /			, ,							,	•	-	•	•		•	•		•	•
max max <td>•</td> <td></td> <td>/</td> <td>•</td> <td>-</td> <td>•</td> <td>•</td> <td></td> <td></td> <td>•</td> <td></td> <td>•</td> <td></td>	•											/	•	-	•	•			•		•	
Mathematic Mathematic Mathematic Application Application <												,	•	-	•	•	•		•	•		
Charter Opport Amount Amo		U 1	/		,		-						•	-	•	•	•		-	-		
Spectral Corrent Max Adds remarks Name Na								\bigcirc				,	•	•	•	•			•		•	
Share Area Share A	Vines & Climbers	Native Bramble	Rubus parvifolius	Rosaceae	Woodland, Forest	1 m	White	0			Sun to semi-shade	Moist to dry	•	•	•	•	•	•	•		•	•
Deep Control Moder Lacence Moder Lacence Moder Lacence Moder Lacen	Shrubs / Small	Common Aotus	Aotus ericoides	Fabaceae	Open Forest, Healthland	< 2 m	Yellow & Red				Sun to semi-shade	Dry to moist	•	•	•	•	•					
Note: Note: <th< td=""><td></td><td>,</td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td>/</td><td>•</td><td>•</td><td>•</td><td>•</td><td>•</td><td></td><td></td><td></td><td></td><td></td></th<>		,										/	•	•	•	•	•					
Shade Add Shade Add Finded Mode Add point Add Add point Add Shade Add point Add					,							/	•	-	•	•			•		•	
Shale, And Charle,					,							/	•	-	•	•	•		-			
Shand, Mar.			1 1									,	•	•	•	•		•	•			•
Option / Inst. Open / Sector Sec							-								•						•	
Duble Puble Numbers (Non- Law Control Numbers (Non- Law Control <th< td=""><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td>· · ·</td><td></td><td>•</td><td>•</td><td>•</td><td>-</td><td></td><td>•</td><td>•</td><td>•</td><td>•</td></th<>												· · ·		•	•	•	-		•	•	•	•
Obset/Addum Prody Current Jub Obset/Addum/Sub Open Fork Jernan Renfer L 1 Renf Prody Current Jub See A second Jub Prody Current Jub Prody Current Jub Prody		,	0/					\bigcirc				/	•	•	•	•					•	
Bindler Michigkein Michigkein <th< td=""><td>Shrubs / Medium</td><td>Prickly Currant Bush</td><td>Coprosma quadrifida</td><td></td><td>Open Forest, Riparian, Rainforest</td><td>2–5 m</td><td>Green-Pale Pink</td><td></td><td></td><td></td><td>Sun to semi-shade</td><td>Wet to moist</td><td></td><td>•</td><td>•</td><td>•</td><td></td><td></td><td></td><td></td><td></td><td></td></th<>	Shrubs / Medium	Prickly Currant Bush	Coprosma quadrifida		Open Forest, Riparian, Rainforest	2–5 m	Green-Pale Pink				Sun to semi-shade	Wet to moist		•	•	•						
Bindar Martin Bindar M	Shrubs / Medium	Hop Bitter Pea		Fabaceae	Woodland, Heathland	1–5 m	Red & Yellow				Sun to semi-shade	Dry	•	•	•	•	•					
Bindlar Madlam View Notice Same Dry Notice Same Dry Notice Same Dry Notice Same Dry Notice Dry					,						Sun to semi-shade	/	•		•	•	•					
Binksy Model Pick Pice Pice Lockgerverun wireverwire Mirase Wooden Header Am Mirase Mir		•												-	•	•	•		-	-	-	
Shinks/ Madium Washing					,							/	-	-		•	•		•	•	•	•
Bindlar Model Network Model Code Model Code Proceeding Proceeding Sum Model Code Sum Model Code <th< td=""><td></td><td>,</td><td>, ,</td><td>,</td><td>, ,</td><td></td><td>-</td><td></td><td></td><td></td><td></td><td>/</td><td>-</td><td></td><td>•</td><td>•</td><td></td><td>•</td><td>•</td><td>-</td><td>•</td><td></td></th<>		,	, ,	,	, ,		-					/	-		•	•		•	•	-	•	
Binds Maxim Marginum production Marking Minisord Q Minis File S S Marking Minisord Q Q Q Q		,	1 1 3	,									-		•	•	•	•	•	-	•	
Bindback Madlum Single Juhl Olice To Kindle Astracce Wir Foreger Woodback Astracce Astrac			,					\bigcirc				,	•	•	•	•		•	•	•	•	•
Brinds / Mailum Horal Brandarris Promotion's apart Promo	Shrubs / Medium	Snowy Daisy Bush	Olearia lirata	Asteraceae	Wet Forest, Woodland	< 4 m	White-Yellow	Õ•			Semi-shade	Moist to dry	•	•					•	•	•	
Binds / Medun Christma Euch Probability Medun Probability Medun San Is cerni-shade Meat to dy Image of the second probability o	Shrubs / Medium	Tree Everlasting	Ozothamnus ferrugineus	Asteraceae	Woodland, Shrubland	2–5 m	White	\bigcirc			Sun	Dry	•	•	•				•			•
Bindbar Solution Solution Solution Solution Moit of oil Solution Moit of oil Solution				Rhamnaceae				0					•	•	•	•			•			
Binds/ Media Days Miler Spin data Pach regions												,	-	•	•	•			•			
Shubs / Jarge Cincuran Walle Accor kproze Southendrates, Weinbrein, Skube 1.5 m Piek Value, Net Southendrates, Weinbrein, Skube Southendrate							-					,	-		•			-	•	•		
Britus / Large Waves, News Bursaria, Blockhom Britus / Large Sun to semi-shade Dry Image: Sun to semi-shade <td></td> <td>,</td> <td></td> <td>•</td> <td></td> <td></td> <td>•</td> <td></td> <td>-</td> <td></td> <td>•</td> <td>•</td> <td></td>		,											•			•		-		•	•	
Bruike Hange Motive Hanke Guedrame Monitor Hank	•				-							,	•	•	•	•		-	•	•	•	•
Bruck / Large Swamp Reperbank Meloleusce enclosing Symme Reperbank Meloleusce e								•ŏ				Moist	•	•	•				•		•	
Trees / Small Golden Wattle Acacia pycnantha Fabaceae Woodland 5 m Yelow Small Full sun Maist lo dry •			/ /		Woodland, Heathland	0.5–4.5 m	White-Cream	00			Sun to semi-shade		•	•	•	•	•		•	•	•	•
Trees / Small Prickly Mases Acacia verticillator Fabaceae Woodlond 3 m Yellow Sun Maist orly •<	•			'				00					-	•	•	•		•	•	•	•	•
Trees / Small Coals Banksia Banksia marginada Proteaceae Wacdland, Heathland 10 m Yellow Image / Small Sun Moist odry Image / Small Sun welto moist Image /												,	-		•	•		•	•	•	•	•
Trees / Small Silver Banksia Banksia marginata Proteaceae Grassland, Woodland 5-11 Vellow Sun Wet to moist • <td></td> <td>,</td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td>,</td> <td>-</td> <td></td> <td>•</td> <td>•</td> <td></td> <td>•</td> <td>•</td> <td>•</td> <td>•</td> <td>•</td>		,										,	-		•	•		•	•	•	•	•
Trees / Small Saw Banksia Banksia serrata Proteaceae Coastal Heathland, Open Forest < 15 m Cream-Yellow Image: Small Sun to semi-shade Dry to moist			-									,	-	-	•				•	•	•	•
Trees / Small River Bottlebrush Callistemon sieberi Myrtaceae Woodland, Riparian 4 m Pink A A B A B A B			-		,								-	-	•	•	•			•	•	
Trees / Small Scented Paperbark Melaleuca squarrosa Myrtaceae Forest, Wetland < 10 m White-Cream Image: Cream Sun to semi-shade Dry Image: Cream <					, ,							,	•	-	•	•	-	•	•	•	•	•
Trees / Medium Silver Wattle Acacia dealbata Fabaceae Woodland, Forest 8-20 m Pale Vellow Sun to semi-shade Dry Image Image Sun to semi-shade Dry Image Im				,				00					•	-	•	•		•	•	-		•
Trees / Medium Drooping Sheak Allocasuarina verticillata Casuarina ceete Grassland, Woodland 5-11 Red & Yellow Sun Sun Dry •					,							Dry	•		•	•		•	•	•	•	•
Trees / Medium Bundy Eucalyptus goniocalyx Myrtaceae Woodland, Open Forest <15 m Cream Cream Sun Dry to moist • <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td>0</td> <td></td> <td></td> <td></td> <td>/</td> <td>•</td> <td></td> <td>•</td> <td>•</td> <td></td> <td>•</td> <td>•</td> <td>•</td> <td>•</td> <td>•</td>								0				/	•		•	•		•	•	•	•	•
Trees / Large Red Gum Eucalyptus canaldulensis Myrtaceae Woodland, Open Forest, Riparian < 40 m Cream Image: Common and the c												/	-		•	•	•					
Trees / Large Broad-Leaved Peppermint Eucalyptus dives Myrtaceae Woodland, Open Forest < 20 m Cream < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m < 1 m <				,									-		•	•		-	•	-	•	•
Trees / LargeYellow GumEucalyptus leucoxylonMyrtaceaeWoodland, Forest<25 mCream-PinkIII </td <td></td> <td></td> <td></td> <td>,</td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td>,</td> <td>-</td> <td>-</td> <td>•</td> <td>•</td> <td></td> <td>•</td> <td>•</td> <td>•</td> <td>•</td> <td>•</td>				,								,	-	-	•	•		•	•	•	•	•
Trees / Large Yellow Box Eucalyptus melliodora Myrtaceae Woodland, Forest 15 m White I Sun Moist odry I												,	•	-	•							
Trees / Large Swamp Gum Eucalyptus ovata Myrtaceae Woodland, Open Forest < 20 m Cream Image: Condebark Sun Moist to wet Image: Condebark Image: Condebark Myrtaceae Woodland, Forest < 20 m Cream Image: Condebark Sun Moist to wet Image: Condebark Image: Condebark Myrtaceae Woodland, Forest < 20 m Cream Image: Condebark Sun Moist to dry Image: Condebark Image: Condebark Sun Moist to dry Image: Condebark Image: Condebark Sun Moist to dry Image: Condebark Image: Condebark Image: Condebark Sun Moist to dry Image: Condebark Image: Condebark Image: Condebark Sun Moist to dry Image: Condebark Image: Condebark Image: Condebark Image: Condebark Image: Condebark Sun Moist to dry Image: Condebark			,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	,								· · · ·		-	•	•		-	•	•	•	•
Trees / Large Candlebark Eucalyptus rubida Myrtaceae Woodland, Forest < 35 m Cream Image: Comparison of the co			,,		,							,	•		•	•			•		•	•
Trees / Large Manna Gum Eucalyptus viminalis Myrtaceae Woodland, Forest 25 m White Image: Control of the large Sun Moist to dry Image: Control of the large Image: Contro of the large Image: Control of	*							Ŏ						-		•						
	•		<i>,</i> ,	Myrtaceae	Woodland, Forest	25 m	White	0					•	•	•	•		•	•	•	•	•
Trees / Large Lilly Pilly Syzgium smithii Myrtaceae Wet Forest, Woodland < 20 m Cream Crea	Trees / Large	Lilly Pilly	Syzygium smithii	Myrtaceae	Wet Forest, Woodland	< 20 m	Cream	\bigcirc			Shade to semi-shade	Wet to moist	•	•	•	•			•		•	

Know your pollinators



European honey bees have two pairs of wings and long, segmented antennae. They are daytime-flying and feed on nectar and pollen. They are generalist pollinators and provide the bulk of pollination services for horticulture and crop plants. Honey bees and native bees are both essential to functioning ecosystems and food security in Australia.

Honey bees have become an important part of the Australian landscape. Honey bees live as colonies, and have a long history of coexistence with humans, including in domestic gardens.





Hoverfly [Fgmily Syrohidge] @Koren Betra **Australian native bees** comprise more than 2000 species, which provide essential pollination services. Native bees are generally solitary and live in nests in the ground or in hollow stems, old borer holes and other cracks and crevices, and some have evolved to pollinate particular native flowers through 'buzz pollination'. Although many Australian native bees are generalist foragers, some species have co-evolved with native plants and adapted to be the most effective pollinators of their flowers. Many native plant species, such as *Dianella* and *Grevillea* require specially adapted insects to access their nectar and enable the transfer of pollen to the stigma. Most native bees are solitary, but some species found in northern Australia (*Tetragonula* sp. and *Austroplebeia* sp.) are social bees and are used for commercial pollination of crops like macadamia nuts.

Fly species number up to 30,000 in Australia, and can be identified by having only one pair of flight wings. A second set of wings are modified into club-shaped paddles that allow flies to hover and stabilise their flight. Unlike bees and wasps, many flies (Brachycera) have very small, clubbed antennae at the front of their head. Flies, including blowflies, are often attracted to flowers that smell like carrion. Some flower-flies, have hairy bodies that easily collect pollen while they are feeding. Flies provide a range of services in the garden, including pollination, decomposition and predation.

Hoverflies are a type of fly, distinguishable by their large eyes, short antennae, bright black and yellow abdomen and their hovering flight behaviour. Adult hoverflies are nectar and pollen feeders. Hoverfly larvae feed on pests such as aphids, thrips and leafhoppers and are excellent biocontrol agents.



Beetles have hard outer wings that form their distinctive beetle shape. Their outer wings form a T-shape where they join at the top, unlike bugs where the outer wings make an X- or Y-shape. Some beetles feed on nectar and pollen, usually by crawling over flower surfaces. There are around 30,000 species of beetles in Australia, with many yet to be formally described.



Butterflies have wings covered in tiny scales. They have clubbed antennae and hold their wings upright when at rest. They are day-flying and have long tongues that they can use to feed on nectar in flowers with deep tubes. Butterflies are usually brightly coloured, with approximately 600 species found in Australia.



Moths also have wings covered in tiny scales and tend to be subtle in colour. They have antennae without clubs and hold their wings flat when at rest. They are generally dusk- and night-flying but there are some exceptions: the grapevine moth is a commonly seen day-flying moth. Moths feed on nectar. Australia has a high diversity of moth species, with up to 22,000 species thought to exist across the continent.

Flower forms



Generalist flowers can be pollinated by many different insects and animals. They are typically saucer shaped with many stamens and have a surface that insects can walk on. *Eucalyptus* flowers and daisy flowers are generalist flowers – they can be pollinated by bees, flies, beetles and butterflies.



Specialist flowers have modifications to their shape and size that only let certain pollinators access the nectar and pollen. These flowers might have deep flower tubes or narrow entry points so that only a select group of pollinators can access them. The advantage of specialisation is that pollination is very targeted and efficient, with accurate pollen placement made possible by co-evolution between flowers and insects. The disadvantage is that if the correct pollinator isn't there, the flowers aren't pollinated. Often, nectar is produced at the base of the flower, forcing pollinators to enter the flower fully and in the process, become covered in pollen.

Pollinator rewards

Nectar is a sugary solution, rich in carbohydrates, vitamins and minerals, produced by flowers and sometimes by glands on leaves or stems (called extra-floral nectaries). Nectar is attractive to insects, and provides an immediate energy source needed for tasks such as hunting pest insects, laying eggs in decomposing organic matter, collecting pollen, or parasitising other insects.

Carbohydrates alone don't support everything needed for health and growth, so insects also need pollen.

Pollen is rich in protein, fats and nutrients. Bees are vegetarian, and need to collect pollen to feed their offspring.

Buzz pollination

Some flowers do not produce any nectar; they specifically target pollencollecting bees, and only offer pollen rewards. To limit pollen loss and ensure effective pollination, some plants produce flowers with specialised, tubular anthers, that only open at the tip. To extract pollen, bees use vibrations to 'buzz' the pollen grains out of the pores of these anthers. Many crops are buzz pollinated, including tomatoes, potatoes, eggplants, capsicum, chillies, tomatillo and cranberries.

European honey bees are unable to buzz pollinate flowers, but several native bees, such as the blue-banded bee, teddy bear bee (*Amegilla* sp.) and carpenter bee (*Xylocopa* sp.) are exceptionally good large buzz pollinators, and have evolved to pollinate native plants such as flax lilies (*Dianella* sp.). Many of our smaller, ground nesting bees utilise vibration to help them excavate their burrows, and they also use that skill to buzz pollen from the anthers of native plants.

Planting buzz-pollinated species will encourage populations of buzz pollinators for successful pollination of food crops and ensure seed set in native plants. Many small ground nesting bees also buzz pollinate native flowers.


Nectar feeding

Grevillea flowers and other tubular flowers are often adapted to be successfully pollinated by birds. Pollen is 'presented' on a floral stigma that extends outside the flower. When birds feed on the nectar, pollen is deposited on their beaks or heads. Bees, also attracted to the sugary nectar, crawl into the side of the flower and feed on the nectar without encountering the pollen-laden stigma. The plant doesn't receive the pollination benefit from the insect, but flowers such Grevillea species can be a very useful source of nectar for insects in the cooler months.



Supported by:







sgaonline.org.au



rbg.vic.gov.au

Nurseries

Most of the plants shown in the planting guide will be available at nurseries that have a good stock of native plants. But if your local nursery doesn't stock the plant you're after, ask them to order it in. For a list of nurseries

that stock all the plants shown in the planting guide, plus other useful resources, visit the Wheen Bee Foundation website or scan the QR code.



WheenBeeFoundation.org.au/our-work/powerful-pollinators

Wheen Bee Foundation

Powerful Pollinators Planting Guides are produced by Wheen Bee Foundation. We fund vital strategic research and education initiatives that strengthen bees, improve pollination efficiency, and protect our food security and ecosystem health. Visit the website for more information.

WheenBeeFoundation.org.au

Far left: The spreading flax lily, *Dianella revoluta,* is buzz pollinated.

Left: This European honey bee is 'side-working': feeding on the nectar-rich flowers without coming into contact with the plant's pollen.

Front cover:

 Australian native bee, Lipotriches species. (Photo: Erica Siegel)
 Aerial view over of south-eastern Melbourne suburbs. (Photo: Dreamstime)
 European honey bees, Apis mellifera. (Photo: Kirrily Hughes)



PP-Vic5.2 (2023) Wheen Bee Foundation 2020. Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License. To view a copy of this license, visit http://creativecommons.org/licenses/by-nc-nd/4.0/