Powerful pollinators

Encouraging insect pollinators in farm landscapes



Pollinators are an essential component of agricultural production and of healthy, biodiverse landscapes. Protecting and enhancing pollinator resources on farms will help support a diverse range of pollinators. This brochure provides an introduction to encouraging insect pollinators on farms, including a guide to choosing plants that will support diverse pollinators throughout the year.



The power of pollinators

Pollinators – mostly insects, but also birds and mammals – assist the formation of seeds and fruit in many plant species by visiting flowers in search of food (nectar and/or pollen). Whilst foraging they transfer pollen from one flower to another, facilitating fertilization, which results in fruits and seeds.

Honey bees, native bees and other native insects like hoverflies, wasps and butterflies provide essential pollination services for native plants, garden flowers, fruits and vegetables.

Pollinators and food security

Without insect pollinators, the quantity and diversity of food and flowers grown in backyard gardens would be severely restricted. Many of the foods we eat, from gardens and farms, benefit from pollination.

Pollinator-dependent foods include citrus, apples, stone-fruits, zucchini, pumpkins, strawberries and tomatoes, as well as plants grown for seed such as sunflowers, coriander and parsley.

The quantity and diversity of insect pollinators are key drivers of production as they influence both food yields and quality. Under-pollination results in smaller and misshapen fruit or seed that isn't viable to grow.

A diverse and healthy community of pollinators generally provides more effective and consistent pollination than relying on any single species.

Pollinators are essential to, and dependent upon, healthy ecosystems. A growing human population and increasing demand for food puts pressure on ecosystems, with potential negative impacts on biodiversity, the environment and food production.



Native vegetation supports pollinators by providing food and nesting sites. Nearby crops and pastures will benefit from the increased abundance and diversity of pollinators in the landscape.

Insect populations are in decline worldwide due to land clearing, intensive or monocultural agriculture, pesticide use, pollution, colony disease, increased urbanisation and climate change. Low pollinator numbers mean not all flowers are pollinated, leading to low fruit or seed set. This in turn reduces fruit and vegetable harvest yields, and decreases food supply.



Under-pollination results in smaller, misshapen fruit such as this strawberry.

Backyard biodiversity

Insect pollinators are a prime example of the importance of healthy ecosystems in urban gardens, parks and reserves. Insects are the 'canaries in the coal mine' of our urban and rural environments. Without our 'littlest creatures', we lack pollinators, natural beneficial pest control services, and critical food source for other insects, birds, amphibians, reptiles and mammals.

The presence of connected and widespread pollinator habitat is critical to support insect populations if we are to maintain sustainable cities and productive, healthy gardens and urban farms for food security and biodiversity.

Pollinators require habitat that contains year-round food sources, breeding resources and nesting sites. The presence of pollinator habitat adjacent to food crops has been shown to improve food production by enabling a greater variety and number of pollinators to persist year-round, providing pollination services when required.

Turn to the centre of this brochure for a guide to planting for pollinators.

Diapause or diet? Where are the insects?

Many insect pollinators undergo a diapause during colder winter months. Diapause is a period of suspended development during unfavourable environmental conditions, and during this period insect pollinators do not need flowers. Birds and other small mammals will continue to benefit from available pollen and nectar during this time.

If there are low numbers of insect pollinators in your local area, it is important to determine whether this is because of diapause, or because of an inadequate availability of nectar and pollen creating a 'food desert' where insect pollinators cannot survive.

There are still many unknowns about insect pollinators in Australia. Take part in Australian Pollinator Week or in the bi-annual Wild Pollinator Count to learn more about pollinators in your area – visit **AustralianPollinatorWeek.org.au** and **WildPollinatorCount.com**

Encouraging pollinators in your garden

Create pollination reservoirs

Pollination reservoirs are areas that provide floral resources for pollinators. They can be gardens, new planting or existing habitat such as established trees, or even local bushland, parks or reserves. A high diversity of plant species is essential to provide nectar, pollen and nesting sites throughout the year. Pollination reservoirs need to be close enough to where pollinators live to ensure that they can fly easily to them.

Improve on what you have

Enhance and improve your existing pollinator habitat where possible. Gardens that already contain established trees, rockeries, ponds, bare soil and organic matter, and a variety of flowering plants, are a valuable resource for beneficial insects and pollinators.

Nature-strips, verges, laneways, vegetable gardens, orchards, nature reserves, and riverbanks and creeks can all be important pollinator-attracting areas. Protect and enhance native pollinator plants in your garden and surrounds for the future.

Plant trees, shrubs and groundcovers

Planting a variety of species of groundcovers, shrubs and trees to in your garden will further attract pollinators to your patch. Initial watering and protection will improve the success rate of young plants. Some species such as wildflowers or native pea species are excellent pollinator attractors and reward careful attention by keen gardeners.

Be a citizen scientist and do some detective work to discover local pollinators in your patch. Visit **inaturalist.ala.org.au** to be involved.

Construct insect real estate

Insect hotels, which are both functional and attractive, are a great way to add to habitat and nesting places for pollinators and insects in your backyard or garden. The hotels are easily moved to be close to flowering plants and those needing pollination, especially if you have a new garden that is still growing. Include lots of different sized holes, cracks and crevices to provide homes for various solitary insect pollinators.

Plant for the future

When establishing pollinator habitat, consider including species that are indigenous to your area but can tolerate increasingly drier and warmer conditions, to create resilient habitat for climate change. Rehabilitate weedy areas into managed pollination reservoirs by introducing lots of flowering plant diversity. Be careful not to plant invasive or listed weeds, and look for suitable replacements.

Amplify the flower signal

Plants have evolved large flowers or clusters of smaller flowers because they attract more pollinator visits. Large, colourful and diverse plantings attract more pollinators. Ideally, plant in groups that contain different vegetation layers – combine a species-rich mixture of wildflowers, ground-covers, herbs, lilies, rushes, climbers, shrubs and trees.

Connectivity counts

Insect pollinators benefit from greater connectivity of habitat in a landscape, which allows them to forage over a wider radius and increase in numbers in a local area. Encourage friends and neighbours to plant for pollinators and create connections in your community.

Get to know your local flora

Your local government area has distinct populations of insects, depending on the local flora and environment. Knowing your local insect species will help you develop better plantings.

The plants growing in nearby nature reserves or bushland will be suited to your climate and soils. Local environment groups and specialist native nurseries can provide information about local plants.

Grow a bumper crop

Pollinator-attracting plants include many fruits and vegetables grown in backyards, community and market gardens, and orchards. Pollinators ensure good yields of crops such as apples, beans, avocado, and almonds, and bush foods such as lilly-pilly and yam daisy.

Reduce chemical use

Insecticides, fungicides and herbicides all affect bee, colony and wild pollinator health. Herbicides can impact pollinators by reducing the availability and diversity of flora and removing vegetation that helps support insect life. Some herbicides can also harm the beneficial bacteria in the insect gut. Insecticides are an obvious threat to pollinators, yet many pollinators will, in healthy numbers, help control pest insects, ultimately reducing the need for insecticide use.

Many crops are dependent on pollination by bees. When chemical pest control is unavoidable, select products that are least harmful for pollinators and apply insecticides in the evening or at night when pollinators are not active. Always use according to directions, especially for withholding periods, and notify beekeepers a few days before spraying chemicals so beehives can be safely relocated away from harm.

Safeguard the bees? The best way to 'save the bees' and protect our pollinators is to create an abundance of diverse habitat — from the ground up! There is much interest in keeping a bee hive to promote pollinators, but there are serious legal and biosecurity responsibilities that must be considered, and that the introduction of a bee hive does not displace existing native pollinators and insects. Be a friend of pollinators and say it with flowers!

A guide to planting for pollinators for Kangaroo Island, South Australia



Healthy populations of insect pollinators are important for sustainable and resilient farms, orchards, gardens, and native flora.

This Guide will help you select plant species to attract and sustain pollinators in agricultural areas and gardens throughout the year.

Kangaroo Island (KI) covers 4430km² with average rainfall of 470-800mm/year mainly in winter. Low-lying lagoonspotted plains of the east rise to ironstone plateau across the centre and west. KI is dominated by primary production and large blocks of native vegetation. The eastern end, and the south and west coasts, are dominated by mallee. Higher rainfall areas in the west support stringybark and sugar gum woodlands. Sheep grazing forms the bulk of KI's primary production, with a lesser amount of cropping and a small amount of horticulture.

The plants listed in this Guide will help supply rewards to pollinators, with an emphasis on species that are indigenous and suited to local climates.

Garden centres sell many common pollinator-attracting ornamental flowers and herbs labelled as 'bee-friendly'.

The eucalypt species in this Guide are mostly large trees, and not suitable for all local environment groups. If you can't gardens, but have been included for their source these plants at your local garden value as good nectar producing species. centre, or indigenous nursery, ask them Most eucalypts do not flower every year, to contact the local wholesale nursery so choosing diverse species will help create continuously flowering habitat.



The pollinator plant list

To create pollinator-attracting plantings, use the Guide to choose a selection of plants with a variety of flower colours, different growth habits and a range of flowering seasons.

For each species, the planting Guide lists:

- life-form/'habit' (climber, herb, shrub or tree) and height (m).
- the vegetation type in which they naturally occur
- flower colour and flowering season
- growth requirements (sun/shade, moist/dry)
- insect groups that may visit each plant and the floral reward (pollen and/or nectar).

The coloured bars indicate the flowering months for each species. Darker shading denotes the peak flowering period, with a lighter shading for non-peak flowering months. Flowering dates may differ between regions and seasons, particularly for non-peak times, if your local climate is consistently warmer or cooler than average, with earlier or later flowering.

Sourcing plants

Most of the plant species listed are available from retail or wholesale nurseries or native plant growers, and suppliers and plant growers listed online. See the reverse of the Guide for details.

WheenBeeFoundation.org.au

			- "				Flowering			Po	linator rewa	rd	Visita	tion by pol	linator	
Lifeform	Common name	Scientific name	Family	Vegetation type	Height	Flower colour	Jan Feb Mar Apr May Jun Jul	Aug Sep Oct Nov Dec	Aspect	Soil moisture	ollen Necto	ar Native bees Hor				Moths Beetles Flies
Crop plants																
Climber / Shrub	Grape Vine	Vitis vinifera	Vitaceae		1–3 m	Green			Sun	Damp to dry	•		• •			•
Forb Forb	Clover Subterranean Clover	Trifolium repens Trifolium subterraneum	Leguminosae		0.3 m 0.3 m	White White-Pink			Sun Sun	Damp to dry Damp to dry	• •	•	•		•	• •
Forb	Onion	Allium cepa	Leguminosae Alliaceae		1m	White			Sun	Damp to dry			•		•	• • •
Forb	Garlic	Allium sativum	Alliaceae		0.5 m	White			Sun	Damp to dry	• •	•	• •	•	•	• • •
Forb	Canola	Brassica napus	Brassicaceae		1.5 m	Yellow			Sun	Damp to dry	• •	•	• •	•		•
Forb	Buckwheat	Fagopyrum esculentum	Polygonaceae		2 m	White			Sun	Damp to dry	•	•	• •	•	•	• •
Forb	Lucerne	Medicago sativa	Leguminosae		0.8 m	Purple			Sun	Damp	• •	•	•		•	• •
Forb	Broad Beans	Vicia faba	Leguminosae		1m	White & Brown			Sun	Damp	• •	•	•		•	• •
Forb Forb	Lupins	Lupinus albus	Leguminosae	Broadacre Cropping	0.5 m 0.5 m	White-Pale Mauve 🔾			Sun	Damp Damp to day	• •	•	•		•	• •
Shrub / Small Tree	Potato Tree Lucerne	Solanum tuberosum Chamaecytisus palmensis	Solanaceae Leguminosae	11 3	2–4 m	Cream			Sun Sun to semi-shade	Damp to dry Damp to dry						• •
Indigenous plants		chamaecynsas paimensis	Leguminosde	rodder / Shener	2-4111	Credin			Sull to settil-situde	Dump to dry		•	•			•
Climber	Purple Coral-Pea	Hardenbergia violacea	Leguminosae	Woodland, Forest, Mallee	1–3 m	Purple			Sun to semi-shade	Damp to dry	• •	•	• •			•
Climber	Old Man's Beard	Clematis microphylla	Ranunculaceae	Woodland, Forest, Mallee	0.5–4 m	Cream			Sun to semi-shade	Damp to dry	•	•	• •			• •
Forb	Chocolate Lily	Arthropodium strictum	Liliaceae	Woodland, Forest	0.3 m	Pink-Mauve			Sun	Damp to dry	•* •	•				
Forb	Black Anther Flax Lily	Dianella revoluta	Liliaceae		1 m	Indigo			Sun to semi-shade	Bamp to ary	•*	•				
Forb	Fairy Fan Flower	Scaevola aemula	Goodeniaceae	Woodland, Forest	0.2 m	Mauve-Purple			Sun	= =	• •	•	• •	•	•	• • •
Forb	Rough Candles	Stackhousia aspericocca	Asteraceae	Woodland, Mallee	0.3 m	Cream-Yellow			Sun to semi-shade	Damp Damp	• •	•	• •	•	•	• • •
Forb Forb	Morning Flag Golden Pennants	Orthrosanthus multiflorus Glischrocaryon behrii	Liliaceae Haloragaceae	Woodland, Mallee. Shrubland Woodland, Mallee, Heathland	0.2–0.5 m 0.3 m	Indigo Yellow			Sun to semi-shade Sun	Damp to dry Damp to dry	•	•	• •			
Forb	Kangaroo Island Pink Eyes	Tetratheca insularis	Elaeocarpeceae		0.3 m	Pink			Sun to semi-shade	Damp to dry	•*		• •	-		
Forb	Native Pelargonium	Pelargonium australe	Geraniaceae	Woodland, Mallee	0.2 m	Pink-Mauve			Sun	Damp to dry	• •		•			
Forb	Coast Swainson Pea	Swainsona lessertiifolia	Leguminosae	Woodland, Mallee, Coastal	0.5 m	Purple			Sun	Dry	• •	•	• •	•		• •
Forb	Leek Lily	Bulbine semibarbata	Liliaceae	Woodland, Mallee	0.2 m	Yellow			Sun to semi-shade	- /	•*	•	•	-		
Forb	Bidgee Widgee	Acaena novae-zelandiae	Rosaceae	Woodland, Mallee	0.4 m	Cream			Sun to semi-shade	Damp to dry	• •	•	•			• •
Forb	Grass Trigger Plant	Stylidium graminifolium	Stylidiaceae	Woodland, Forest, Mallee	0.3–0.5 m	Pink			Sun to semi-shade	Damp to dry	•	•	• •	•		•
Sedge	Sticky Sword Sedge	Lepidospemum viscidum	Cyperaceae	Woodland, Forest, Heathland, Mallee, Wetland		White-Cream			Sun to semi-shade	Damp to dry	• •	•	• •	•		• • •
Sedge	Tussock Sedge	Carex appressa	Cyperaceae	Wetland	1–1.7 m	Yellow-Brown			Sun	Damp to wet	•	•		•		• •
Sedge	Saw Sedge	Gahnia trifida	Cyperaceae		1–2 m	Brown			Sun	Wet to damp	• •	•	•	•	•	•
Shrub	Creeping Boobialla	Myoporum parvifolium	Myoporaceae	Shrubland, Coastal	0.3 m	White Cream-Green			Sun to semi-shade Sun to semi-shade	Damp to dry	• •	•	• •	•	•	• • •
Shrub Shrub	Shiny Ground Berry Grass Tree	Acrotriche patula Xanthorrea semiplana semiplana	Epacridaceae Liliaceae	Woodlsnd, Mallee Woodland, Forest, Mallee, Riverine	0.3–1.5 m 1-3 m	Cream			Sun to semi-shade	Dry Dry			•			• • •
Shrub	Muntries	Kunzea pomifera	Myrtaceae	Woodland, Mallee, Coastal	0.3 m	White			Sun	Dry			• •			
Shrub	Coastal Beard Heath	Leaucopogon parviflorus	Apecridaceae	Woodland, Shrubland, Coastal	0.5–2.5 m	White			Sun	Damp to dry	• •	•	• •	•	•	• • •
Shrub	Scented Groundsel	Senecio odoratus	Asteraceae	Woodland, Mallee, Shrubland	0.5–1.8 m	Yellow			Sun to semi-shade	Damp to dry	• •	•	• •	•	•	• • •
Shrub	Leafless Bitter Pea	Daviesia brevifolia	Leguminosae	Woodland, Mallee, Heathland	0.3–0.5 m	Red & Orange 🛛 🔴			Sun	Dry	• •	•	• •			•
Shrub	Kangaroo Island Bitter Pea	Daviesia asperula	Leguminosae		0.5–1.5 m	Brown & Orange 🛛 🔵			Sun to semi-shade	Dry	• •	•	• •			•
Shrub	Hop Goodenia	Goodenia ovata	Goodeniaceae		0.4–0.8 m				Sun to semi-shade	Damp to dry	• •	•	•			•
Shrub	Mat Heath Myrtle	Thryptomene ericaea	Myrtaceae	Woodland, Forest, Mallee	0.5–2 m	White-Pink			Sun to semi-shade	Dry	• •	•	• •	•	•	• • •
Shrub	Common Fringe Myrtle	Calytrix tetragona	Myrtaceae	Woodland, Heathland, Mallee	0.5–1.7 m	White-Pink			Sun	Dry	• •	•	• •	•	•	• • •
Shrub Shrub	Smooth Fringe Myrtle Round Leaf Kangaroo Island Correa	Calytrix glaberrima Correa backhouseana var orbicularis	Myrtaceae	Woodland, Heathland, Mallee Woodland, Mallee, Shrubkand, Coastal	0.5–1.5 m	White-Pink O			Sun Sun to semi-shade	Dry Dry	• •		• •		•	
Shrub	Kangaroo Island Correa	Correa reflexa var insularis	Rutaceae		1–2.5 m	Pale Green			Sun to semi-shade	Dry			•	•		
Shrub	Wrinkled Hakea	Hakea rugosa	Proteaceae	Woodland, Mallee, Wetland	0.5–1.7 m	White			Sun	Damp to dry	• •	•	• •	•	•	• • •
Shrub	Grey Trymalium	Tyrmalium wayi	Rhamnaceae	Woodland, Mallee	1–3 m	Pale Yellow			Sun	Dry	• •	•	•			• •
Shrub	Small Darwinia	Darwinia micropetala	Myrtaceae	Woodland, Mallee, Heathland, Wetland	0.4–1.5 m	White-Pink			Sun	Damp to dry	• •	•	• •	•	•	• • •
Shrub	Coast Bitter Bush	Adriana quadripartita	Euphorbiaceae	Woodland, Mallee, Shrubland, Coastal	1–3 m	Red-Brown & Yellow-Green 🛑 🤇			Sun	Damp to dry	• •	•	• •	•	•	• • •
Shrub	Spiny Mint Bush	Prostanthera spinosa	Labiatae	Woodland, Forest	0.5–1 m	Pale Mauve			Sun to semi-shade	Damp to dry	• •	•	•	•		•
Shrub	Heath Tea Tree	Leptospermum myrsinoides	Myrtaceae		1–3 m	White-Pink			Sun	2.)	• •	•	• •	•	•	• • •
Shrub / Small Tree	Prickly Moses	Acacia verticillata	Leguminosae	Woodland, Forest, Wetland	2–3 m	Yellow			Sun	Damp to dry	•	•	• •	•		• • •
Shrub / Small Tree Shrub / Small Tree	Scarlet Bottlebrush Prickly Tea Tree	Callistemon rugulosus Leptospermum continentale	Myrtaceae Myrtaceae	Woodland, Mallee, Wetland Riverine, Wetland, Heathland, Forest	1.5–4 m 1–4 m	Red White-Pink			Sun Sun	Damp Damp		•	• •	-	•	• • •
Shrub / Small Tree	Woolly Tea Tree	Leptospermum lanigerum	Myrtaceae	Riverine, Woodland, Heathland	1–4 m	White			Sun to semi-shade	Wet to damp			• •			• • •
Shrub / Small Tree	Dtyland Tea Tree	Melaleuca lanceolata	Myrtaceae	Woodland, Forest, Mallee	2–6 m	White			Sun to semi-shade	Dry	• •	•	• •	•	•	
Shrub / Small Tree	Slender Honey Myrtle	Melaleuca gibbosa	Myrtaceae	Woodland, Forest, Mallee, Wetland, Riverine		Pink			Sun to semi-shade	Damp to dry	• •	•	• •	•	•	• • •
Shrub / Small Tree	Broom Honey Myrtle	Melaleuca uncinata	Myrtaceae		2–5 m	Cream			Sun to semi-shade	Damp to dry	•	•	• •			• • •
Shrub / Small Tree	Christmas Bush	Bursaria spinosa	Pittosporaceae		2–6 m	White-Cream			Sun to semi-shade	Damp	• •	•	• •	•	•	• • •
Shrub / Small Tree	Silver Banksia	Banksia marginata	Proteaceae		2–5 m	Pale Yellow			Sun to semi-shade	Damp to dry	• •	•	• •	•		• • •
Shrub / Small Tree	Beaked Hakea	Hakea rostrata	Proteaceae	Woodland, Heathland, Mallee	1–5 m	White			Sun to semi-shade	Dry		•	• •	•	•	• • •
Shrub / Small Tree	Desert Hakea	Hakea mitchellii	Proteaceae		1–4 m	White (Sun	Dry		•	• •	•		• • •
Shrub / Small Tree Tree	Native Juniper Drooping Sheoak	Myoporum insulare Allocasuarina verticillata	Myoporaceae Casuarinaceae		1–5 m 3–10 m	Red & Yellow			Sun Sun	Dry Dry		•	• •	•	•	• • •
Tree	Swamp Wattle	Anocasuarina verncinara Acacia provincialis	Leguminosae	, ,	3–10 m 3–8 m	Pale Yellow			Sun to semi-shade	- '	•		• •	•		
Tree	Coastal Wattle	Acacia leiopylla	Leguminosae		2–3 m	Yellow			Sun	Dry	•		• •	•		• • •
Tree	Golden Wattle	Acacia pycnantha	Leguminosae	Woodland, Forest, Mallee	2–5 m	Yellow			Sun to semi-shade	Damp to dry	•	•	• •	•		• • •
Tree	Kingscote Mallee	Eucalyptus rugosa	Myrtaceae		3–20 m	White			Sun	Dry	• •	•	• •		•	• • •
Tree	Kangaroo Island Narrow Leaf Mallee	Eucalyptus cneorifolia	, Myrtaceae	Woodland, Forest, Mallee	3–15 m	White			Sun	Damp to dry	• •	•	• •	•	•	• • •
Tree	Brown Stringybark	Eucalyptus baxteri	Myrtaceae	Woodland, Forest	20 m	White-Cream			Sun	Damp to dry	• •	•	• •		•	• • •
Tree	River Red Gum	Eucalyptus camaldulensis	Myrtaceae	Riverine, Forest	40 m	White			Sun	= =	• •	•	• •	•	•	• • •
Tree	SA Blue Gum	Eucalyptus leucoxylon	Myrtaceae	Riverine, Woodland, Forest	30 m	Cream-Pink			Sun	Damp to dry	• •	•	• •	•	•	• • •
Tree	Pink Gum	Eucalyptus fasciculosa	Myrtaceae	Woodland, Forest, Mallee, Riverine	8–25 m				Sun to semi-shade	Damp to dry		•	• •	•	•	• • •
	Cup Gum Coastal White Mallee	Eucalyptus cosmophylla	Myrtaceae		2–6 m 3–10m	White (Sun to semi-shade Sun	Damp to dry	• •		• •	•		• • •
Tree Tree	Messmate Stringybark	Eucalyptus diversifolia Eucalyptus obliqua	Myrtaceae Myrtaceae	Woodland, Mallee, Coastal Woodland, Forest	3–10m 20 m	White-Cream			Sun	Damp to dry Dry						
Tree	Post Lincoln Mallee	Eucalyptus obliqua Eucalyptus albopurpurea	Myrtaceae	Woodland, Mallee	4–20 m	White-Pink			Sun to semi-shade	Damp to dry		•	• •			• • •
Tree	Sugar Gum	Eucalyptus cladocalyx	Myrtaceae	Riverine, Woodland, Forest	40 m	White			Sun	Dry	• •	•	•	•	•	• • •
Tree	Kangaroo Island Mallee Ash	Eucalyptus remota	Myrtaceae		2–10 m	White-Cream			Sun	- '	• •	•	• •	•	•	• • •
	-															

Know your pollinators



European honey bees have two pairs of wings and long, segmented antennae. They are daytime-flying and feed on nectar and pollen. They are generalist pollinators and provide the bulk of pollination services for horticulture and crop plants. Honey bees and native bees are both essential to functioning ecosystems and food security in Australia.

Honey bees have become an important part of the Australian landscape. Honey bees live as colonies, and have a long history of coexistence with humans, including in domestic gardens.







Australian native bees comprise more than 2000 species, which provide essential pollination services. Native bees are generally solitary and live in nests in the ground or in hollow stems, old borer holes and other cracks and crevices, and some have evolved to pollinate particular native flowers through 'buzz pollination'. Although many Australian native bees are generalist foragers, some species have co-evolved with native plants and adapted to be the most effective pollinators of their flowers. Many native plant species, such as *Dianella* and *Grevillea* require specially adapted insects to access their nectar and enable the transfer of pollen to the stigma. Most native bees are solitary, but some species found in northern Australia (*Tetragonula* sp. and *Austroplebeia* sp.) are social bees and are used for commercial pollination of crops like macadamia nuts.

Fly species number up to 30,000 in Australia, and can be identified by having only one pair of flight wings. A second set of wings are modified into club-shaped paddles that allow flies to hover and stabilise their flight. Unlike bees and wasps, they have very small, clubbed antennae at the front of their head. Flies, including blowflies, are often attracted to flowers that small like carrion; they generally have hairy bodies that easily collect pollen while they are feeding. Flies provide a range of services in the garden, including pollination, decomposition and predation.

Hoverflies are a type of fly, distinguishable by their large eyes, short antennae, bright black and yellow abdomen and their hovering flight behaviour. Adult hoverflies are nectar and pollen feeders. Hoverfly larvae feed on pests such as aphids, thrips and leafhoppers and are useful biocontrol agents.



Beetles have hard outer wings that form their distinctive beetle shape. Their outer wings form a T-shape where they join at the top, unlike bugs where the outer wings make an X- or Y-shape. Beetles feed on nectar and pollen, usually by crawling over flower surfaces. There are around 30,000 species of beetles in Australia, with many yet to be formally described.



Butterflies have wings covered in tiny scales. They have clubbed antennae and hold their wings upright when at rest. They are day-flying and have long tongues that they can use to feed on nectar in flowers with deep tubes. Butterflies are usually brightly coloured, with approximately 600 species found in Australia.



Moths also have wings covered in tiny scales and tend to be subtle in colour. They have antennae without clubs and hold their wings flat when at rest. They are generally dusk- and night-flying but there are some exceptions: the grapevine moth is a commonly seen day-flying moth. Moths feed on nectar. Australia has a high diversity of moth species, with up to 22,000 species thought to exist across the continent.

Flower forms



Generalist flowers can be pollinated by many different insects and animals. They are typically saucer shaped with many stamens and have a surface that insects can walk on. *Eucalyptus* flowers and daisy flowers are generalist flowers – they can be pollinated by bees, flies, beetles and butterflies.



Specialist flowers have modifications to their shape and size that only let certain pollinators access the nectar and pollen. These flowers might have deep flower tubes or narrow entry points so that only a select group of pollinators can access them. The advantage of specialisation is that pollination is very targeted and efficient, with accurate pollen placement made possible by co-evolution between flowers and insects. The disadvantage is that if the correct pollinator isn't there, the flowers aren't pollinated. Often, nectar is produced at the base of the flower, forcing pollinators to enter the flower fully and in the process, become covered in pollen.

Pollinator rewards

Nectar is a sugary solution, rich in carbohydrates, vitamins and minerals, produced by flowers and sometimes by glands on leaves or stems (called extra-floral nectaries). Nectar is attractive to insects, and provides an immediate energy source needed for tasks such as hunting pest insects, laying eggs in decomposing organic matter, collecting pollen, or parasitising other insects.

Carbohydrates alone don't support everything needed for health and growth, so insects also need pollen.

Pollen is rich in protein, fats and nutrients. Bees are vegetarian, and need to collect pollen to feed their offspring.

Buzz pollination

Some flowers do not produce any nectar; they specifically target pollencollecting bees, and only offer pollen rewards. To limit pollen loss and ensure effective pollination, some plants produce flowers with specialised, tubular anthers, that only open at the tip. To extract pollen, bees use vibrations to 'buzz' the pollen grains out of the pores of these anthers. Many crops are buzz pollinated, including tomatoes, potatoes, eggplants, capsicum, chillies, tomatillo and cranberries.

European honey bees are unable to buzz pollinate flowers, but several native bees, such as the blue-banded bee, and teddy bear bee (Amegilla sp.) and carpenter bee (Xylocopa sp.) are exceptionally good large buzz pollinators, and have evolved to pollinate native plants such as flax lilies (Dianella sp.). Many of our smaller, ground nesting bees utilise vibration to help them excavate their burrows, and they also

use that skill to buzz pollen from the anthers of native plants.

Planting buzz-pollinated species will encourage populations of buzz pollinators for successful pollination of food crops and ensure seed set in native plants. Many small ground nesting bees also buzz pollinate native flowers.

Nectar feeding

Grevillea flowers and other tubular flowers are often adapted to be successfully pollinated by birds. Pollen is 'presented' on a floral stigma that extends outside the flower. When birds feed on the nectar, pollen is deposited on their beaks or heads. Bees, also attracted to the sugary nectar, crawl into the side of the flower and feed on the nectar without encountering the pollen-laden stigma. The plant doesn't receive the pollination benefit from the insect, but flowers such Grevillea species can be a very useful source of nectar for insects in the cooler months.



Supported by:



dess.net.au



adelaide.edu.au



samuseum.sa.gov.au

Wholesale Nurseries

Most of the plants shown in the planting guide will be available at nurseries that have a good stock of native plants. But if your local nursery doesn't stock the plant you're after, ask them to order it in. For a list of wholesale nurseries

that stock all the plants shown in the planting guide, plus other useful resources, visit the Wheen Bee Foundation website or scan the QR code.



WheenBeeFoundation.org.au/our-work/powerful-pollinators

Wheen Bee Foundation

Powerful Pollinators Planting Guides are produced by Wheen Bee Foundation. We fund vital strategic research and education initiatives that strengthen bees, improve pollination efficiency, and protect our food security and ecosystem health. Visit the website for more information.

WheenBeeFoundation.org.au

Far left: The spreading flax lily, Dianella revoluta, is buzz pollinated.

Left: This European honey bee is 'side-working': feeding on the nectar-rich flowers without coming into contact with the plant's pollen.

Front cover:

Xylocopa aerata. (Photo: Remko Leijs) 2. Sunrise, Mouth Flat, Kangaroo Island. (Photo: Richard Glatz) 3. European honey bees, Apis mellifera. (Photo: Kirrily Hughes)





anz.com.au

frrr.org.au

PP-SA13.1 (2021) Wheen Bee Foundation 2021. Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License. To view a copy of this license, visit http://creativecommons.org/licenses/by-nc-nd/4.0/