Powerful pollinators

Encouraging insect pollinators in farm landscapes



Pollinators are an essential component of agricultural production and of healthy, biodiverse landscapes. Protecting and enhancing pollinator resources on farms will help support a diverse range of pollinators. This brochure provides an introduction to encouraging insect pollinators on farms, including a guide to choosing plants that will support diverse pollinators throughout the year.



The power of pollinators

Pollinators – mostly insects, but also birds and mammals – assist the production of seeds and fruit in many plant species by visiting flowers in search of food (nectar and/or pollen). Whilst foraging they transfer pollen from one flower to another, facilitating fertilisation, which results in fruits and seeds.

Honey bees, native bees and other native insects like hoverflies, wasps and butterflies provide essential pollination services for native plants, garden flowers, fruits and vegetables.

Pollinators and food security


Without insect pollinators, the quantity and diversity of food and plants grown in backyard gardens would be severely restricted. Many of the foods we eat, from gardens and farms, benefit from pollination.

Pollinator-dependent foods include citrus, apples, stone-fruits, zucchini, pumpkins, strawberries and tomatoes, as well as plants grown for seed such as sunflowers, coriander and parsley.

The quantity and diversity of insect pollinators are key drivers of production as they influence both food yields and quality. Under-pollination results in smaller and misshapen fruit or seed that isn't viable.

A diverse and healthy community of pollinators generally provides more effective and consistent pollination than relying on any single species.

Pollinators are essential to, and dependent upon, healthy ecosystems. A growing human population and increasing demand for food puts pressure on ecosystems, with potential negative impacts on biodiversity, the environment and food production.



Native vegetation supports pollinators by providing food and nesting sites. Nearby crops and pastures will benefit from the increased abundance and diversity of pollinators in the landscape.

Insect populations are in decline worldwide due to land clearing, intensive or monocultural agriculture, pesticide use, pollution, colony disease, increased urbanisation and climate change. Low pollinator numbers mean not all flowers are pollinated, leading to low fruit or seed set. This in turn reduces fruit and vegetable harvest yields, and decreases food supply.



Under-pollination results in smaller, misshapen fruit such as this strawberry.

Backyard biodiversity

Insect pollinators are a prime example of the importance of healthy ecosystems in urban gardens, parks and reserves. Insects are the 'canaries in the coal mine' of our urban and rural environments. Without our 'littlest creatures', we lack pollinators, natural beneficial pest control services, and critical food source for other insects, birds, amphibians, reptiles and mammals.

The presence of connected and widespread pollinator habitat is critical to support insect populations if we are to maintain sustainable cities and productive, healthy gardens and urban farms for food security and biodiversity.

Pollinators require habitat that contains year-round food sources, breeding resources and nesting sites. The presence of pollinator habitat adjacent to food crops has been shown to improve food production by enabling a greater variety and number of pollinators to persist year-round, providing pollination services when required.

Turn to the centre of this brochure for a guide to planting for pollinators.

Diapause or diet? Where are the insects?

Many insect pollinators undergo a diapause during colder winter months. Diapause is a period of suspended development during unfavourable environmental conditions, and during this period insect pollinators do not need flowers. Birds and other small mammals will continue to benefit from available pollen and nectar during this time.

If there are low numbers of insect pollinators in your local area, it is important to determine whether this is because of diapause, or because of an inadequate availability of nectar and pollen creating a 'food desert' where insect pollinators cannot survive.

There are still many unknowns about insect pollinators in Australia. Take part in Australian Pollinator Week or in the annual Australian Pollinator Count to learn more about pollinators in your area – visit: **AustralianPollinatorWeek.org.au** and **AustralianPollinatorCount.au**

Encouraging pollinators in your garden

Create pollination reservoirs

Pollination reservoirs are areas that provide floral resources for pollinators. They can be gardens, new plantings or existing habitat such as established trees, or even local bushland, parks or reserves. A high diversity of plant species is essential to provide nectar, pollen and nesting sites throughout the year. Pollination reservoirs need to be close enough to where pollinators live to ensure that they can fly easily to them.

Improve on what you have

Enhance and improve your existing pollinator habitat where possible. Gardens that already contain established trees, rockeries, ponds, bare soil and organic matter, and a variety of flowering plants, are a valuable resource for beneficial insects and pollinators.

Nature-strips, verges, laneways, vegetable gardens, orchards, nature reserves, and riverbanks and creeks can all be important pollinator-attracting areas. Protect and enhance native pollinator plants in your garden and surrounds for the future.

Plant trees, shrubs and groundcovers

Planting a variety of species of groundcovers, shrubs and trees in your garden will further attract pollinators to your patch. Initial watering and protection will improve the success rate of young plants. Some plants such as wildflowers or native pea species are excellent at attracting pollinators, rewarding keen gardeners with a diversity of native pollinators.

Be a citizen scientist and do some detective work to discover local pollinators in your patch. Visit **inaturalist.ala.org.au** to be involved.

Construct insect real estate

Insect hotels, which are both functional and attractive, are a great way to add to habitat and nesting places for pollinators and insects in your backyard or garden. The hotels are easily moved to be close to flowering plants and those needing pollination, especially if you have a new garden that is still growing. Include lots of different sized holes, cracks and crevices to provide homes for various solitary insect pollinators.

Plant for the future

When establishing pollinator habitat, consider including species that are indigenous to your area but can tolerate increasingly drier and warmer conditions, to create resilient habitat for climate change. Rehabilitate weedy areas into managed pollination reservoirs by introducing lots of flowering plant diversity. Be careful not to plant invasive or listed weeds, and look for suitable replacements.

Amplify the flower signal

Plants have evolved large flowers or clusters of smaller flowers because they attract more pollinator visits. Large, colourful and diverse plantings attract more pollinators. Ideally, plant in groups that contain different vegetation layers – combine a species-rich mixture of wildflowers, ground-covers, herbs, lilies, rushes, climbers, shrubs and trees.

Connectivity counts

Insect pollinators benefit from greater connectivity of habitat in a landscape, which allows them to forage over a wider radius and increase in numbers in a local area. Encourage friends and neighbours to plant for pollinators and create connections in your community.

Get to know your local flora

Your local government area has distinct populations of insects, depending on the local flora and environment. Knowing your local insect species will help you develop better plantings.

The plants growing in nearby nature reserves or bushland will be suited to your climate and soils. Local environment groups and specialist native nurseries can provide information about local plants.

Grow a bumper crop

Pollinator-attracting plants include many fruits and vegetables grown in backyards, community and market gardens, and orchards. Pollinators ensure good yields of crops such as apples, beans, avocado, and almonds, and bush foods such as Lilly-Pilly and Finger Limes.

Reduce chemical use

Insecticides, fungicides and herbicides all affect bee, colony and wild pollinator health. Herbicides can impact pollinators by reducing the availability and diversity of flora and removing vegetation that helps support insect life. Some herbicides can also harm the beneficial microbes in the insect gut. Insecticides are an obvious threat to pollinators, yet many beneficial insects will, in healthy numbers, help control pest insects, ultimately reducing the need for insecticide use.

Many crops are dependent on pollination by bees. When chemical pest control is unavoidable, select products that are least harmful for pollinators and apply insecticides in the evening or at night when pollinators are not active. Always use according to directions, especially for withholding periods, and notify beekeepers a few days before spraying chemicals so beehives can be safely relocated away from harm.

Safeguard the bees? The best way to 'save the bees' and protect our pollinators is to create an abundance of diverse habitat — from the ground up! There is much interest in keeping a bee hive to promote pollinators, but there are serious legal and biosecurity responsibilities that must be considered, and that the introduction of a bee hive does not displace existing native pollinators and insects. Be a friend of pollinators and say it with flowers! A guide to planting for pollinators for Townsville, Queensland

Healthy populations of insect pollinators are important for sustainable and resilient farms, orchards, gardens, and native flora.

This Guide will help you select plant species to attract and sustain pollinators in agricultural areas and gardens throughout the year.

Townsville Local Government Area (LGA) is • the vegetation type in which they environmentally biodiverse, encompassing naturally occur part of the Brigalow Belt Bioregion, Wet Tropics Bioregion and Einasleigh Uplands bioregion. The Brigalow Belt is a shrubby open forest on clay soils; the Wet Tropics encroach the northern portion of the LGA and is characterised by complex sclerophyll forests, woodlands and rainforest; and Einasleigh Uplands is in the west of the LGA and contains Eucalypt forests on hills, ranges and plateaus. The region has dry tropical climate with two distinct seasons: the wet season (December – March) and the dry season (April – November).

The plants listed in this Guide will help supply rewards to pollinators, with an emphasis on species that are indigenous later flowering. and suited to local climates.

Garden centres sell many common pollinator-attracting ornamental flowers Most of the plant species listed are and herbs labelled as 'bee-friendly'.

The eucalypt species in this Guide are mostly large trees, and not suitable for all gardens, but have been included for their source these plants at your local garden value as good nectar producing species. centre, or indigenous nursery, ask them Most eucalypts do not flower every year, to contact the local wholesale nursery so choosing diverse species will help create continuously flowering habitat.





The pollinator plant list

To create pollinator-attracting plantings, use the Guide to choose a selection of plants with a variety of flower colours, different growth habits and a range of flowering seasons.

For each species, the planting Guide lists:

- life-form/'habit' (climber, herb, shrub or tree) and height (m).
- flower colour and flowering season
- growth requirements (sun/shade, moist/dry)
- insect groups that may visit each plant and the floral reward (pollen and/or nectar).

The coloured bars indicate the flowering months for each species. Darker shading denotes the peak flowering period, with a lighter shading for non-peak flowering months. Flowering dates may differ between regions and seasons, particularly for non-peak times, if your local climate is consistently warmer or cooler than average, with earlier or

Sourcing plants

available from retail or wholesale nurseries or native plant growers, and local environment groups. If you can't suppliers and plant growers listed online. See the reverse of the Guide for details.

Lifeform	Common name	Scientific name	Family	Vegetation type	
Indigenous plants					
Forbs / Herbs	Native Ginger	Alpinia caerulea	Zingiberaceae	Sub-Tropical Rainforest	
Forbs / Herbs	Australian Crepe Ginger	Cheilocostus potierae	Costaceae	Rainforest	F
Forbs / Herbs	Common Everlasting	Chrysocephalum apiculatum	Asteraceae	Grassland, Open Woodland	H
Forbs / Herbs	Yellow Button	Coronidium rupicola	Asteraceae	Open, Wet Sclerophyll, Rainforest	F
Forbs / Herbs	Rattlepod	Crotalaria brevis	Fabaceae	Sclerophyll Forest	F
Forbs / Herbs	Spurge	Euphorbia bifida	Euphorbiaceae	Scrub, Forest margins	F
Forbs / Herbs	Small St. John's Wort	Hypericum gramineum	Hypericaceae	Open Forest, Grassland	F
Forbs / Herbs	Birdsville Indigo	Indigofera linnaei	Fabaceae	Wood & Shrubland, Wooded Grassland	F
Forbs / Herbs	Beach Morning Glory	Ipomoea pes-caprae	Convolvulaceae	Coastal Sand Dunes	
Forbs / Herbs	Grass Lily	Murdannia graminea	Commelinaceae	Sclerophyll Forest	Γ
Forbs / Herbs	Pastel Flower	Pseuderanthemum variabile	Acanthaceae	Rainforest, Wet Scleorphyll Forest	
Forbs / Herbs	Arrowroot	Tacca leontopetaloides	Dioscoreaceae	Rainforest, Monsoon Forest	Γ
Forbs / Herbs	Trephrosia	Tephrosia filipes	Fabaceae	Sclerophyll Forest	
Forbs / Herbs	Native Violet	Viola banksii	Violaceae	Coastal, Rainforest fringes	Γ
Forbs / Herbs	Golden Everlasting	Xerochrysum bracteatum	Asteraceae	Forest, Woodland, Grassland	
Forbs / Herbs	Blue Flax-Lily	Dianella caerulea	Asphodelaceae	Sclerophyll Forest	
Grass/Sedge/Rush	Water Chestnut	Eleocharis dulcis	Cyperaceae	Permanent freshwater	
Grass/Sedge/Rush	Many-Flowered Mat-Rush	Lomandra multiflora	Asparagaceae	Forest, Woodland	
Grass/Sedge/Rush	Awned Panick Grass	Oplismenus compositus	Poaceae	Forest	
Grass/Sedge/Rush	Kangaroo Grass	Themeda triandra	Poaceae	Variable	
Lilies & Irises	Stream Lily	Crinum pedunculatum	Amaryllidaceae	Forest, Wetland	
Shrub (small)	Wattle	Acacia umbellata	Fabaceae	Eucalypt Woodland, Rocky slopes & ridges	
Shrub (small)	Great Woolly Malayan Lilac	Callicarpa candicans	Lamiaceae	Rain & Monsoon Forest, Vine Thickets	
Shrub (small)	Blue-Flower Rattlepod	Crotalaria verrucosa	Fabaceae	Open Forest, Riparian, Vine Thickets	
Shrub (small)	Blue Tongue	Melastoma malabathricu	Melastomataceae	Rain, Monsoon & Open Forest, Heath	
Shrub (small)	Pavetta	Pavetta australiensis	Rubiaceae	Rain, Monsoon & Beach Forests	L
Shrub (small)	Phyllanthus	Phyllanthus lamprophyllus	Phyllanthaceae	Open and/or Monsoon Forest	
Shrub (small)	Pink Pea-Bush	Tephrosia brachyodon	Fabaceae	Woodland, Dry Sclerophyll Forest	L
Shrub (large)	Townsville Wattle	Acacia leptostachya	Fabaceae	Woodland, Grassland	-
Shrub (large)	Heathlands Wattle	Acacia simsii	Fabaceae	Eucalypt Open Forest, Woodland	
Shrub (small)	Gardenia White Star	Gardenia psidioides	Rubiaceae	Open Eucalypt & Monsoon Forest	-
Shrub (large)	Hovea	Hovea longipes	Fabaceae	Rainforest, Scrub, Woodland	H
Shrub (large)	Mueller's Evodia	Melicope rubra	Rutaceae	Drier Rainforest	-
Shrub (large) Shrub (large)	Boobialla Brush Muttonwood	Myoporum acuminatum Myrsine variabilis	Scrophulariaceae Primulaceae	Rain, Wet Sclerophyll & Casaurina Forest Rain & Monsoon Forest	H
Shrub (large)	Wild Yellow Jasmine	Pittosporum revolutum	Pittosporaceae	Rain, Wet Sclerophyll & Open Forest	H
Shrub (large)	Beach Naupaka	Scaevola taccada	Goodeniaceae	Sandy Beaches, Rocky outcrops	H
Shrub (large)	Stenocarpus	Stenocarpus angustifolius	Proteaceae	Dry Eucalypt Forest	F
Shrub (large)	Milkwood/Coolaroo	Wrightia saligna	Apocynaceae	Open Forest, Savannah Woodland	F
Shrub (large)	Golden Penda	Xanthostemon chrysanthus	Myrtaceae	Coastal Rainforest	Ē
Tree (small)	Club-Leaf Wattle	Acacia hemignosta	Fabaceae	Open Woodland	F
Tree (small)	Native Gardenia	Atractocarpus fitzalanii	Rubiaceae	Rainforest	Γ
Tree (small)	Lemon-Scented Myrtle	Backhousia citriodora	Myrtaceae	Coastal Rainforest	
Tree (small)	Red Bloodwood	Corymbia erythrophloia	Myrtaceae	Open Eucalyptus Woodland	Γ
Tree (small)	Tuckeroo	Cupaniopsis anacardioides	Sapindaceae	Littoral Rainforest, Scrub	
Tree (small)	Grey Corkwood	Erythrina vespertilio	Fabaceae	Dry Rainforest	Γ
Tree (small)	Mount Stuart Ironbark	Eucalyptus paedoglauca	Myrtaceae	Open Forest, Grassy Woodland	
Tree (small)	Broad-Leaved Ballart	Exocarpos latifolius	Santalaceae	Variable, Open habitats	
Tree (small)	Python Tree	Gossia bidwillii	Myrtaceae	Rainforest	
Tree (small)	Fern-Leaf Grevillea	Grevillea pteridifolia	Proteaceae	Open Forest	
Tree (small)	Canary Beech	Huberantha nitidissima	Annonaceae	Rainforest	
Tree (small)	Kamala	Macaranga tanarius	Euphorbiaceae	Rainforest	L
Tree (small)	Yellow Ball Flower	Mallotus nesophilus	Euphorbiaceae	Beach Forest, Monsoon Forest	
Tree (small)	Fibrebark	Melaleuca nervosa	Myrtaceae	Open Forest, Woodland	Ŀ
Tree (small)	Weeping Bottlebrush	Melaleuca viminalis	Myrtaceae	Rainforest and Open Forest	
Tree (small)	White Cedar	Melia azedarach	Meliaceae	Subtropical and Dry Rainforest	L
Tree (small)	Brown Damson	Terminalia arenicola	Combretaceae	Beach Forest, Closed Forest	F
Tree (small)	Mueller's Damson	Terminalia muelleri	Combretaceae	Beach Forest, Monsoon Forest	-
Tree (large) Tree (large)	White Siris	Albizia procera Alectryon connatus	Fabaceae	Open, Rain and Monsoon Forest	
	Alectryon Illawarra Flame Tree	Brachychiton acerifolius	Sapindaceae Malvaceae	Variable Forests	-
Tree (large)		Corymbia tessellaris	Malvacede	Subtropical Rainforest Woodland, Riparian corridors	-
Tree (large) Tree (large)	Moreton Bay Ash/Carbeen Wild Tamarind	Diploglottis diphyllostegia	Sapindaceae	Rainforest	\vdash
Tree (large)	Grey Ironbark	Eucalyptus crebra	Myrtaceae	Woodland, Open Forest	H
Tree (large)	Forest Red Gum	Eucalyptus tereticornis	Myrtaceae	Wet and Dry Sclerophyll Forest	H
Tree (large)	Small-Leaved Fig	Ficus obliqua	Moraceae	Littoral Rainforest, Subtropical Rainforest	F
Tree (large)	Teak	Flindersia australis	Rutaceae	Dry Rainforest, Subtropical Rainforest	F
Tree (large)	Ribbon Fan Palm	Livistona decora	Arecaceae	Dry & Riparian Rainforest, Open Rorest	F
Tree (large)	Northern Swamp Box	Lophostemon grandiflorus	Myrtaceae	Open Forest, Rainforest margins	
Tree (large)	Paper Bark	Melaleuca fluviatilis	Myrtaceae	Sclerophyll Forest, Swampy Open Forest	f
Tree (large)	Pink-Flowered Doughwood	Melicope elleryana	Rutaceae	Subtropical Rainforest, Swamp Forest	
Tree (large)	Brush Cherry	Syzygium australe	Myrtaceae	Forest	f
Tree (large)	Damson Plum	Terminalia sericocarpa	Combretaceae	Dry Rainforest, Rainforest, Monsoon Forest	
Vines & Climbers	Wombat Berry	Eustrephus latifolius	Asparagaceae	Sclerophyll Forest, Woodland, Heath	ſ
Vines & Climbers	Scrambling Lily	Geitonoplesium cymosum	Asphodelaceae	Sclerophyll Forest, Woodland	
Vines & Climbers	Native Hoya	Hoya australis	Apocynaceae	Rainforest	ſ
Vines & Climbers	Jasmine	Jasminum simplicifolium	Oleaceae	Drier Rainforest, Monsoon Forest	
Vines & Climbers	Wonga Wonga Vine	Pandorea pandorana	Bignoniaceae	Forest, Woodland	ſ
Vines & Climbers	Giant Pepper Vine	Piper hederaceum	Piperaceae	Rainforest	
Vines & Climbers	Climbing Cassia	Senna gaudichaudii	Fabaceae	Open Eucalypt, Rain & Monsoon Forest	
	č			-

WheenBeeFoundation.org.au

		Flowering	Flowering		Coll moliature	Pollinator reward				Visitation by pollinator					
Height	Flower colour	Jan Feb Mar Apr May Jun Jul Aug Sep Oct No	ov Dec	Aspect	Soil moisture	Pollen	Nectar	Native bees	Honey bees	Hoverflies	Wasps	Butterflies	Moths	Beetles	Flies
2–2.5 m	White			Sun to full shade	Moist	•	•	٠	•			•			
< 4 m	Red-Cream () Yellow			Semi-shade to shade Full sun		•		•		•		•		•	•
0.4 m	Yellow			Sun to semi-shade	Moist to dry Dry to moist	•	•	•	•	•	•	•	•	•	•
0.3 m	Yellow			Sun to semi-shade		•	•	•	•			•			
0.3 m	White Yellow			Full sun to light shade		•	•	•	•	•	•	•		•	•
0.1–0.4 m 0.15–0.5 m	Red			Sun to semi-shade Sun to semi-shade	Well-drained Dry, well-drained	•	•	•	•	•	•	•		•	•
Creeping	Pink			Full sun	Well-drained	•	•	•	•	•	•	•	•	•	•
0.1–0.6 m	Pink		- 22	Sun to semi-shade Sun to light shade	Well-drained Moist, well-drained	•	•	•	•	•		•			
0.15-0.3 m	White, Mauve			J	Moist, well-drained	•	•	•				•		•	•
< 2 m	Purple			Full sun	Dry, well-drained	•	•	•	•			•			
0.2–0.4 m	, ,			Light to heavy shade	. ,	•	•	•	•	•	•			_	
0.2–0.8 m <1 m				Full sun Sun to semi-shade	Moist to dry Well-drained	• •*	•	•	•	•	•	•		•	•
< 1.5 m	White-brown			Full sun	Wet	•								•	•
0.2-0.9 m	Cream				Variable	•	•	•	•		•			•	•
0.3–0.8 m < 1.2 m	Pink Green, Brown		100	Semi-shade to shade Sun to semi-shade	Moist Variable	•								•	•
0.3–2 m	White			Sun to semi-shade		•	•	•		•			•	•	
es < 3 m	Golden			Full sun	Dry, well-drained	•	•	•	•	•	•	•		•	•
2 m <1 m	Purple Purple			Sun to semi-shade Full sun	Dry to moderate Dry to moist	•	•	•	•	•	•	•	•	•	•
1–3 m	Pink			Full sun to full shade	,	•*		•	•						
1–5 m	Cream			Full sun to full shade	Moist	•	•	•	•	•	•	•	•	•	•
1–2 m	Cream			Sun to semi-shade	Dry to moderate	•	•						•		•
1–2 m 2–5 m	Pink Yellow			Sun to semi-shade Full sun	Dry, well-drained Well drained	•	•	•	•	•	•			•	•
2–4 m	Golden			Full sun	Dry, well-drained	•		•	•	•	•	•		•	•
1 m	White			Sun to semi-shade	Well-drained	•	•					•	•		
< 5 m	Purple Pink			Sun to semi-shade Sun to semi-shade	Variable	•	•	•	•	•	•	•	•	•	•
t < 13 m	Cream, Purple			Sun to light-shade	Well-drained	•	•	•	•	•	•	•	•	•	•
3–12 m	Cream			Semi to full shade	Moist to moderate	•	•	•	•	•	•	•	•	•	•
< 3 m	Yellow			Sun to semi-shade	Moist to moderate	•	•	•	•	•	•	•	•	•	•
< 4 m 4–5 m	Cream		-	Sun to light shade Sun to light shade	Well-drained Dry, well-drained	•	•	•	•		•	•		•	
2–3 m	Cream-yellow			Full sun	Dry, well-drained	•	•	•	•	•	•			•	•
10–15 m	Golden			Full sun	Moist to moderate	•	•	•	•	•	•	•		•	•
3–10 m < 6 m	Golden Cream			Full sun Sun to half-shade	Dry, well-drained Moist to moderate	•	•	•	•	•	•	•		•	•
< 8 m	Cream		- 10	Sun to half-shade	Moist to moderate	•	•	•	•	•	•	•	•	•	•
< 8 m	Cream			Full sun	Dry to moderate	•	•	٠	•	•	•	•	•	•	•
<12 m	Cream, Green, Pink 🤇			Full sun	Moderate, well-drained	•	•	•	•	•	•	•		•	•
5–12 m ~10 m	Red, Purple			Full sun Full sun	Moist to moderate Dry, well-drained	•	•	•	•	•	•	•		•	•
2–10 m					Dry, well-drained	•		•	•	•	•	•		•	•
18 m	White			Semi to full shade	Moist to moderate	•	•	•	•	•	•	•		•	•
4–8 m <10 m	Golden Cream-yellow		i	Full sun Sun to semi-shade	Dry, well-drained	•	•	•	•		•			•	
<10 m	Cream-yellow		-	Sun to semi-shade		•	•	•	•	•	•			•	•
1.5–6 m	Cream-yellow			Sun to semi-shade		•	•	•	•	•	•			•	•
4–10 m 2–10 m	Cream (Red			Sun to semi-shade Sun to semi-shade		•		•	•	•	•	•	•	•	•
<12 m	Cream, Pink			Sun to semi-shade		•	•	•	•	•	•	•	-	•	•
<10 m	White			Sun to semi-shade	Well-drained	•	•	•	•	•	•			•	•
6-8 m	White Cream-yellow			Sun to semi-shade Sun to light-shade		•	•	•	•	•	•			•	•
< 25 m 10–20 m				Sun to light-shade Sun to half-shade	Variable, well drained	•	•	•	•	•	•	•	•	•	•
< 35 m	Red			Sun to semi-shade	Moist to moderate	•	•	•	•	•	•	•	-	•	•
< 35 m	Cream			Sun to semi-shade		•	•	•	•	•	•	•	•	•	•
< 30 m < 35 m	Cream Cream			Semi to full shade Sun to semi-shade	Moist to moderate	•	•	•	•	•	•	•	•	•	•
< 50 m	Cream			Sun to semi-shade		•	•	•	•	•	•	•	•	•	•
st < 30 m	Inconspicuous			Sun to semi-shade	/	•					•				
< 40 m	White Cream-yellow			Sun to semi-shade Sun to semi-shade	Variable, well drained	•	•	•	•	•	•	•		•	•
< 10 m	Cream			Full sun	Dry to moderate	•	•	•	•	•	•	•	•	•	•
t < 30 m	Cream-yellow			Sun to semi-shade	Variable	•	•	•	•	•	•	•	•	•	•
< 25 m	Pink			Sun to semi-shade		•	•	•	•	•	•	•		•	•
< 35 m st 10-40 m	White (White			Sun to semi-shade Sun to semi-shade	Moist Variable, well drained	•	•	•	•	•	•	•	•	•	•
Climber	Cream, Pink 🛛 🔾			Sun to semi-shade		•	•					•	•	•	•
Climber	White			Semi to full shade	Dry to moist	•	•	٠	•	•				•	•
< 6 m Climber	Cream (White			Semi to full shade Sun to semi-shade	Well-drained	•	•	•	•	•	•	•	•	•	•
> 6 m				Sun to semi-shade		•	•	•	•	•	•		•	•	•
Climber	Cream, Green			Semi to full shade	Moist, well-drained	•									•
2–3 m	Yellow			Sun to semi-shade	Well-drained	•*		•	•						

Know your pollinators



European honey bees have two pairs of wings and long, segmented antennae. They are daytime-flying and feed on nectar and pollen. They are generalist pollinators and provide the bulk of pollination services for horticulture and crop plants. Honey bees and native bees are both essential to functioning ecosystems and food security in Australia.

Honey bees have become an important part of the Australian landscape. Honey bees live as colonies, and have a long history of coexistence with humans, including in domestic gardens.





Hoverfly [Family Syrophidae] © Karen Betra **Australian native bees** comprise more than 2000 species, which provide essential pollination services. Native bees are generally solitary and live in nests in the ground or in hollow stems, old borer holes and other cracks and crevices, and some have evolved to pollinate particular native flowers through 'buzz pollination'. Although many Australian native bees are generalist foragers, some species have co-evolved with native plants and adapted to be the most effective pollinators of their flowers. Many native plant species, such as *Dianella* and *Grevillea* require specially adapted insects to access their nectar and enable the transfer of pollen to the stigma. Most native bees are solitary, but some species found in northern Australia (*Tetragonula* sp. and *Austroplebeia* sp.) are social bees and are used for commercial pollination of crops like macadamia nuts.

Fly species number up to 30,000 in Australia, and can be identified by having only one pair of flight wings. A second set of wings are modified into club-shaped paddles that allow flies to hover and stabilise their flight. Unlike bees and wasps, many flies (Brachycera) have very small, clubbed antennae at the front of their head. Flies, including blowflies, are often attracted to flowers that smell like carrion. Some flower-flies, have hairy bodies that easily collect pollen while they are feeding. Flies provide a range of services in the garden, including pollination, decomposition and predation.

Hoverflies are a type of fly, distinguishable by their large eyes, short antennae, bright black and yellow abdomen and their hovering flight behaviour. Adult hoverflies are nectar and pollen feeders. Hoverfly larvae feed on pests such as aphids, thrips and leafhoppers and are excellent biocontrol agents.



Beetles have hard outer wings that form their distinctive beetle shape. Their outer wings form a T-shape where they join at the top, unlike bugs where the outer wings make an X- or Y-shape. Some beetles feed on nectar and pollen, usually by crawling over flower surfaces. There are around 30,000 species of beetles in Australia, with many yet to be formally described.



Butterflies have wings covered in tiny scales. They have clubbed antennae and hold their wings upright when at rest. They are day-flying and have long tongues that they can use to feed on nectar in flowers with deep tubes. Butterflies are usually brightly coloured, with approximately 600 species found in Australia.



Moths also have wings covered in tiny scales and tend to be subtle in colour. They have antennae without clubs and hold their wings flat when at rest. They are generally dusk- and night-flying but there are some exceptions: the grapevine moth is a commonly seen day-flying moth. Moths feed on nectar. Australia has a high diversity of moth species, with up to 22,000 species thought to exist across the continent.

Flower forms



Generalist flowers can be pollinated by many different insects and animals. They are typically saucer shaped with many stamens and have a surface that insects can walk on. *Eucalyptus* flowers and daisy flowers are generalist flowers – they can be pollinated by bees, flies, beetles and butterflies.



Specialist flowers have modifications to their shape and size that only let certain pollinators access the nectar and pollen. These flowers might have deep flower tubes or narrow entry points so that only a select group of pollinators can access them. The advantage of specialisation is that pollination is very targeted and efficient, with accurate pollen placement made possible by co-evolution between flowers and insects. The disadvantage is that if the correct pollinator isn't there, the flowers aren't pollinated. Often, nectar is produced at the base of the flower, forcing pollinators to enter the flower fully and in the process, become covered in pollen.

Pollinator rewards

Nectar is a sugary solution, rich in carbohydrates, vitamins and minerals, produced by flowers and sometimes by glands on leaves or stems (called extra-floral nectaries). Nectar is attractive to insects, and provides an immediate energy source needed for tasks such as hunting pest insects, laying eggs in decomposing organic matter, collecting pollen, or parasitising other insects.

Carbohydrates alone don't support everything needed for health and growth, so insects also need pollen.

Pollen is rich in protein, fats and nutrients. Bees are vegetarian, and need to collect pollen to feed their offspring.

Buzz pollination

Some flowers do not produce any nectar; they specifically target pollencollecting bees, and only offer pollen rewards. To limit pollen loss and ensure effective pollination, some plants produce flowers with specialised, tubular anthers, that only open at the tip. To extract pollen, bees use vibrations to 'buzz' the pollen grains out of the pores of these anthers. Many crops are buzz pollinated, including tomatoes, potatoes, eggplants, capsicum, chillies, tomatillo and cranberries.

European honey bees are unable to buzz pollinate flowers, but several native bees, such as the blue-banded bee, and teddy bear bee (*Amegilla* sp.) and carpenter bee (*Xylocopa* sp.) are exceptionally good large buzz pollinators, and have evolved to pollinate native plants such as flax lilies (*Dianella* sp.). Many of our smaller, ground nesting bees utilise vibration to help them excavate their burrows, and they also use that skill to buzz pollen from the anthers of native plants.

Planting buzz-pollinated species will encourage populations of buzz pollinators for successful pollination of food crops and ensure seed set in native plants. Many small ground nesting bees also buzz pollinate native flowers.

Nectar feeding

Grevillea flowers and other tubular flowers are often adapted to be successfully pollinated by birds. Pollen is 'presented' on a floral stigma that extends outside the flower. When birds feed on the nectar, pollen is deposited on their beaks or heads. Bees, also attracted to the sugary nectar, crawl into the side of the flower and feed on the nectar without encountering the pollen-laden stigma. The plant doesn't receive the pollination benefit from the insect, but flowers such Grevillea species can be a very useful source of nectar for insects in the cooler months.



Nurseries

Most of the plants shown in the planting guide will be available at nurseries that have a good stock of native plants. But if your local nursery doesn't stock the plant you're after, ask them to order it in. For a list of wholesale nurseries

that stock all the plants shown in the planting guide, plus other useful resources, visit the Wheen Bee Foundation website or scan the QR code.



WheenBeeFoundation.org.au/our-work/powerful-pollinators

Wheen Bee Foundation

Powerful Pollinators Planting Guides are produced by Wheen Bee Foundation. We fund vital strategic research and education initiatives that strengthen bees, improve pollination efficiency, and protect our food security and ecosystem health. Visit the website for more information.

WheenBeeFoundation.org.au

Far left: The spreading flax lily, *Dianella revoluta,* is buzz pollinated.

Left: This European honey bee is 'side-working': feeding on the nectar-rich flowers without coming into contact with the plant's pollen.

Front cover:

 Amegilla sp. on Melastoma sp. (Photo: Laura Lopresti)
 Paluma Range National Park. (Photo: Alejandro de la Fuente)
 European honey bees, Apis mellifera. (Photo: Kirrily Hughes)

Supported by:



townsville.qld.gov.au



jcu.edu.au

PP-QLD24.1 (2023) Wheen Bee Foundation 2023. Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License. To view a copy of this license, visit http://creativecommons.org/licenses/by-nc-nd/4.0/

